arXiv:1201.4129v1 [math.ST] 19 Jan 2012

Theoretical Results on FIEGARCH Processes

Sílvia R.C. Lopes* and Taiane S. Prass
Mathematics Institute - UFRGS
Porto Alegre - RS - Brazil

January 20, 2012

Abstract

Here we present a theoretical study on the main properties of FIEGARCH processes. We analyze the conditions for the existence, the invertibility, the stationarity and the ergodicity of these processes. We prove that, if $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH(p,d,q) process then, under mild conditions, $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA$(q,d,0)$ process. The convergence order for the polynomial coefficients that describes the volatility is presented and results related to the spectral representation and to the covariance structure of both processes $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ are also discussed. Expressions for the kurtosis and the asymmetry measures for any stationary FIEGARCH(p,d,q) process are also derived. The h-step ahead forecast for the processes $\{X_t\}_{t\in\mathbb{Z}}$, $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is given with their respective mean square error forecast. The work also presents a Monte Carlo simulation study showing how to generate, estimate and forecast based on five different FIEGARCH models. The forecasting performance of ARCH-type models is compared through an empirical application to Brazilian stock market exchange index.

Keywords. Long-Range Dependence, Volatility, Stationarity, Ergodicity, FIEGARCH Processes.
Mathematics Subject Classification (2000). Primary 60G10, 62G05, 62G35, 62M10, 62M15; Secondary 62M20.

1 Introduction

Financial time series present an important characteristic known as volatility which can be defined/measured in different ways but it is not directly observable. A common approach, but not unique, is to define the volatility as the conditional standard deviation (or the conditional variance) of the process and use heteroskedastic models to describe it.

ARCH-type models, proposed by Engle (1982), constitute one of the main classes of econometric models used for representing the dynamic evolution of volatilities. Another popular one is the class of Stochastic Volatility (SV) models. A drawback of this class is that the volatility is specified as a latent variable which is not directly observable and this can make the estimation challenging. By ARCH-type models we mean not only the ARCH(p) model proposed by Engle (1982), which characterizes the volatility as a function of powers of past observed values (consequently, the volatility

---

*Corresponding author. E-mail: silvia.lopes@ufrgs.br

can be observed one-step ahead), but also the several generalizations that were lately proposed to properly model the dynamics of the volatility.

Among the generalizations of the ARCH model are the so-called Fractionally Integrated Exponential GARCH (FIEGARCH), introduced by Bollerslev and Mikkelsen (1996), and Fractionally Integrated GARCH (FIGARCH), proposed by Baillie et al. (1996). These models generalize, respectively, the EGARCH model (Nelson, 1991), and the GARCH model (Engle and Bollerslev, 1986), and they capture the effects of long-memory characteristic in the conditional variance (see Doukhan et al., 2003).

FIEGARCH models have not only the capability of modeling clusters of volatility (as in the ARCH and GARCH models) and capturing its asymmetry[1] (as in the EGARCH models) but they also take into account the characteristic of long memory in the volatility (as in the FIGARCH models, with the advantage of been weakly stationary if $d < 0.5$). Besides non-stationarity (in the weak sense), another drawback of the FIGARCH(p, d, q) models is that we must have $d \geq 0$ and the polynomial coefficients in its definition must satisfy some restrictions so the conditional variance will be positive. FIEGARCH(p, d, q) models do not have this problem since the variance is defined in terms of the logarithm function.

Some authors argue that the long memory behavior observed in the sample autocorrelation and periodogram functions of financial time series could actually be caused by the non-stationarity property. According to Mikosch and Stărică (1999), long range behavior could be just an artifact due to structural changes. On the other hand, Mikosch and Stărică (1999) also argue that, when modeling return series with large sample size, considering a single GARCH model is unfeasible and that the best alternative would be to update the parameter values along the time. As an alternative to the traditional heteroskedastic models, Ming (2000) presents a regime switching model that, combined with heavy tailed distributions, presents the long memory characteristic.

It is our belief that FIEGARCH models are a competitive alternative for modeling large sample sized data, especially because it avoids parameter updating. Also, as we prove in this work, FIEGARCH processes are weakly stationary if and only if $d < 0.5$ and hence, non-stationarity can be easily identified. Moreover, Saadi et al. (2006) analyze the daily returns of the Tunisian stock market and rule out the random walk hypothesis. According to the authors, the rejection of this hypothesis seems to be due to substantial non-linear dependence and not to non-stationarity in the return series and, after comparing several ARCH-type models they concluded that a stationary FIEGARCH model provides the best fit for the data. Furthermore, Jayasuriya (2009) presents a sub period investigation of long memory and structural changes in volatility. The authors consider FIEGARCH models to examine the long run persistence of stock return volatility for 23 developing markets for the period of January 2000 to October 2007. No clear evidence that long memory characteristic could be attributed to structural changes in volatility was found.

In practice, usually a simple FIEGARCH(p, d, q) model with $p, q \in \{0, 1\}$ suffices to fully describe the data. For instance, Saadi et al. (2006) and Ruiz and Veiga (2008), consider FIEGARCH$(1, d, 0)$ models while Jayasuriya (2009) considers FIEGARCH$(1, d, 1)$ models. However, for the time series considered in the working papers by Prass and Lopes (2011a, 2011b), higher values of p and q are needed to provide a good fit for the data. Thus, in this paper we present the main theoretical properties of FIEGARCH(p, d, q) processes, for any $p, q \geq 0$, extending the results known in the literature for the particular cases $p, q \in \{0, 1\}$. We also present a simulated study including generation, estimation and forecasting features of FIEGARCH models. Knowing the theoretical properties of these processes is fundamental since any estimator used to fit a model to a time series will assume some properties such as stationarity and ergodicity. On the other hand, working with simulated data is important in order to analyze the finite sample behavior of the estimators.

---

[1] By asymmetry we mean that the volatility reacts in an asymmetrical form to the returns, that is, volatility tends to rise in response to “bad” news and to fall in response to “good” news.

Although different definitions of FIEGARCH processes can be found in the literature (see, for instance, Zivot and Wang, 2005 and Palma, 2001), we consider in this work the formulation proposed by Bollerslev and Mikkelsen (1996), which depends on the function $\lambda(z) = \frac{\alpha(z)}{\beta(z)}(1-z)^{-d}$. The conditions for the existence of power series representation for this function and the behavior of the coefficients in this representation are investigated throughout this work. This study is fundamental not only for simulation purposes but also to draw conclusions on the autocorrelation and spectral density functions decay of the non-observable process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and the observable one $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$. We also provide a recurrence formula to calculate the coefficients of this series expansion for any $p, q \geq 0$. This recurrence formula allows to easily simulate FIEGARCH processes.

We consider the fact that, given a FIEGARCH(p,d,q) process $\{X_t\}_{t\in\mathbb{Z}}$, it is immediate that $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA(q,d,p) process, and we analyze the existence of an ARFIMA representation for the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$. This is very useful since the parameter estimation for ARFIMA models is well known (see Lopes, 2008 and references therein) and, to the best of our knowledge, this result is absent in the literature. Another contribution of this work is to provide the expressions for the asymmetry and kurtosis measures of FIEGARCH(p,d,q) process, for all $p, q \geq 0$. This result extends the particular case considered in Ruiz and Veiga (2008). The ARFIMA representation of $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and the result that any FIEGARCH process is a martingale difference with respect to the natural filtration $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$, where $\mathcal{F}_t := \sigma(\{Z_s\}_{s\leq t})$, are applied to obtain the h-step ahead forecast for the processes $\{X_t\}_{t\in\mathbb{Z}}$ and $\{X_t^2\}_{t\in\mathbb{Z}}$. We also present the h-step ahead forecast for both $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ processes, with their respective mean square error forecast. To the best of our knowledge, formal proves for these expressions are not given in the literature of FIEGARCH(p,d,q) processes.

Despite the fact that the quasi-likelihood is one of the most applied method in non-linear process estimation, asymptotic results for FIEGARCH process are still an open question (see Palma, 2007)[2]. Therefore, we consider here a simulated study to investigate its finite sample performance. Since it is expected that, the better the fit, the better the forecasting, we also investigate the fitted models' forecasting performance.

The paper is organized as follows: Section 2 presents the formal definition of FIEGARCH process and its theoretical properties. We give a recurrence formula to obtain the coefficients in the power series expansion of the polynomial that describes the volatility and we show their asymptotic properties. The autocovariance and spectral density functions of the processes $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ are also presented and analyzed. The asymmetry and kurtosis measures of any stationary FIEGARCH process are also presented. Section 3 presents the theoretical results regarding the forecasting. Section 4 presents a Monte Carlo simulation study including the generation of FIEGARCH time series, estimation of the model parameters and the forecasting based on the fitted model. Section 5 presents the analysis of an observed time series and the comparison of the forecasting performance for different ARCH-type models. Section 6 concludes the paper.

2 FIEGARCH Process

In this section we present the *Fractionally Integrated Exponential Generalized Autoregressive Conditional Heteroskedastic* process (FIEGARCH). This class of processes, introduced by Bollerslev and

---

[2] The asymptotic properties for the quasi-likelihood method are well established for ARCH/GARCH models (see, for instance, Lee and Hansen, 1994; Lumsdaine, 1996; Berkes et al., 2003; Berkes and Horváth, 2003 and Hall and Yao, 2003) and also for EGARCH models (see, for instance, Straumann and Mikosch, 2003).

Mikkelsen (1996), describes not only the volatility varying on time and the volatility clusters (known as ARCH/GARCH effects) but also the volatility long-range dependence and its asymmetry.

Here, we present some results related to the existence, stationarity and ergodicity for these processes. We analyze the autocorrelation and the spectral density functions decay for both $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ processes. Conditions for the existence of a series expansion for the polynomial that describes the volatility are given and a recurrence formula to calculate the coefficients of this expansion is presented. We also discuss the coefficients asymptotic behavior. We observe that if $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH(p,d,q) process then, $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA(q,d,p) process and we prove that, under mild conditions, $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA$(q,d,0)$ process with correlated innovations. We present the expression for the kurtosis and the asymmetry measures for any stationary FIEGARCH(p,d,q) process.

Throughout the paper, given $a \in \mathbb{R}\cup\{-\infty,+\infty\}$, $f(x) = O(g(x))$ means that $|f(x)| \leq c|g(x)|$, for some $c > 0$, as $x \to a$; $f(x) = o(g(x))$ means that $f(x)/g(x) \to 0$, as $x \to a$; $f(x) \sim g(x)$ means that $f(x)/g(x) \to 1$, as $x \to a$. We also say that $f(x) \approx g(x)$, as $x \to \infty$, if for any $\varepsilon > 0$, there exists $x_0 \in \mathbb{R}$ such that $|f(x) - g(x)| < \varepsilon$, for all $x \geq x_0$. Also, given any set T, T^* corresponds to the set $T\backslash\{0\}$ and $\mathbb{I}_A(\cdot)$ is the indicator function defined as $\mathbb{I}_A(z) = 1$, if $z \in A$, and 0, otherwise.

From now on, let $(1-\mathcal{B})^d$ be the operator defined by its Maclaurin series expansion as,

$$(1-\mathcal{B})^d = \sum_{k=0}^{\infty} \frac{\Gamma(k-d)}{\Gamma(k+1)\Gamma(d)} := \sum_{k=0}^{\infty} \delta_{d,k}\,\mathcal{B}^k := \delta_d(\mathcal{B}), \tag{2.1}$$

where $\Gamma(\cdot)$ is the gamma function, $\mathcal{B}$ is the backward shift operator defined by $\mathcal{B}^k(X_t) = X_{t-k}$, for all $k \in \mathbb{N}$, and the coefficients $\delta_{d,k}$ are such that $\delta_{d,0} = 1$ and $\delta_{d,k-1} = \delta_{d,k-1}\left(\frac{k-1-d}{k}\right)$, for all $k \geq 1$.

Remark 2.1. Note that expression (2.1) is valid only for non-integer values of d. When $d \in \mathbb{N}$, $(1-\mathcal{B})^d$ is merely the difference operator $1-\mathcal{B}$ iterated d times. Also, one observe that, upon replacing d by $-d$, the operator $(1-\mathcal{B})^{-d}$ has the same binomial expansion as the polynomial given in (2.1), that is

$$(1-\mathcal{B})^{-d} = \sum_{j=0}^{\infty} \delta_{-d,j}\mathcal{B}^j := \sum_{k=0}^{\infty} \pi_{d,k}\mathcal{B}^k, \tag{2.2}$$

where $\pi_{d,j} = \delta_{-d,j}$, for all $j \in \mathbb{N}$. Moreover, $\pi_{d,k} \sim \frac{1}{\Gamma(d)\,k^{1-d}}$, as $k \to \infty$ (see Prass, 2008). Therefore, $\pi_{d,k} = O(k^{d-1})$, as k goes to infinity.

Suppose that $\{Z_t\}_{t\in\mathbb{Z}}$ is a sequence of independent and identically distributed (i.i.d.) random variables, with zero mean and variance equal to one. Let $\alpha(\cdot)$ and $\beta(\cdot)$ be the polynomials of order p and q defined, respectively, by

$$\alpha(z) = \sum_{i=0}^{p}(-\alpha_i)z^i = 1 - \sum_{i=1}^{p}\alpha_i z^i \quad \text{and} \quad \beta(z) = \sum_{j=0}^{q}(-\beta_j)z^j = 1 - \sum_{j=1}^{q}\beta_j z^j, \tag{2.3}$$

with $\alpha_0 = \beta_0 = -1$. We assume that $\beta(z) \neq 0$, if $|z| \leq 1$, and that $\alpha(\cdot)$ and $\beta(\cdot)$ have no common roots. These conditions assure that the operator $\frac{\alpha(\mathcal{B})}{\beta(\mathcal{B})}$ is well defined.

Definition 2.1. The process $\{X_t\}_{t\in\mathbb{Z}}$ is a *Fractionally Integrated* EGARCH *process*, denoted by FIEGARCH(p,d,q), if

$$X_t = \sigma_t Z_t, \tag{2.4}$$

$$\ln(\sigma_t^2) = \omega + \frac{\alpha(\mathcal{B})}{\beta(\mathcal{B})}(1-\mathcal{B})^{-d}g(Z_{t-1}), \quad \text{for all } t \in \mathbb{Z}, \tag{2.5}$$

where $\omega \in \mathbb{R}$ and $g(\cdot)$ is defined by

$$g(Z_t) = \theta Z_t + \gamma[|Z_t| - \mathbb{E}(|Z_t|)], \quad \text{for all } t \in \mathbb{Z}, \text{ with } \theta, \gamma \in \mathbb{R}. \tag{2.6}$$

Remark 2.2. Note that, in Definition 2.1, no conditions on the parameter d are imposed. Necessary and sufficient conditions on the parameter d, to guarantee the existence of the stochastic process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$, satisfying (2.5), are discussed in the sequel. Also notice that when $d = 0$, we obtain the well known EGARCH process. Moreover, the theory presented here can be easily adapted to a more general definition where one may consider

$$\ln(\sigma_t^2) = \omega + \sum_{k=0}^{\infty} c_k g(Z_{t-1}), \quad \text{for all } t \in \mathbb{Z},$$

where $\{c_k\}_{k\in\mathbb{N}}$ is a sequence of real numbers for which the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is well defined, $\{Z_t\}_{t\in\mathbb{Z}}$ is white noise process with variance not necessarily equal to one and $g(\cdot)$ is any measurable function.

Figure 2.1 (a) presents a simulated time series $\{X_t\}_{t=1}^n$, with $n = 2,000$ observations, derived from a FIEGARCH$(0, d, 1)$ process. Figure 2.1 (b) and (c) present the conditional variance $\{\sigma_t^2\}_{t=1}^n$ and the logarithm of the conditional variance $\{\ln(\sigma_t^2)\}_{t=1}^n$, both related to the same time series as in Figure 2.1 (a). For this simulation we set $d = 0.3578$, $\theta = -0.1661$, $\gamma = 0.2792$, $\omega = -7.2247$ and $\beta_1 = 0.6860$. These are the parameter values of the FIEGARCH model fitted to the Bovespa index log-returns in Prass and Lopes (2011b).



(a) $\{X_t\}_{t=1}^{2000}$ (b) $\{\sigma_t^2\}_{t=1}^{2000}$ (c) $\{\ln(\sigma_t^2)\}_{t=1}^{2000}$

Figure 2.1: (a) Simulated FIEGARCH time series $\{X_t\}_{t=1}^n$, with $n = 2,000$ observations. (b) The conditional variance of $\{X_t\}_{t=1}^n$ and (c) the logarithm of the conditional variance.

Notice that the function $g(\cdot)$, given by the expression (2.6), can be rewritten as

$$g(Z_t) = \begin{cases} (\theta + \gamma)Z_t - \gamma\mathbb{E}(|Z_t|), & Z_t \geq 0; \\ (\theta - \gamma)Z_t - \gamma\mathbb{E}(|Z_t|), & Z_t < 0. \end{cases}$$

This expression clearly shows the asymmetry in response to positive and negative returns. Also, it is easy to see that $g(\cdot)$ is non-linear if $\theta \neq 0$ and the asymmetry is due to the values of $\theta \pm \gamma$. While the parameter θ, also known in the literature as *leverage parameter*, shows the return's sign effect, the parameter γ denotes the return's magnitude effect. Therefore, the model is able to capture the fact that a negative return usually results in higher volatility than a positive one.

From Prass (2008) it is known that if $\{Z_t\}_{t\in\mathbb{Z}}$ is a sequence of independent and identically distributed random variables with zero mean and variance equal to one, then $\{g(Z_t)\}_{t\in\mathbb{Z}}$ is a strictly stationary and ergodic process. Moreover, if $\mathbb{E}(Z_t^2) < \infty$, it is also weakly stationary (consequently, it is a white noise process) with mean zero and variance σ_g^2, given by

$$\sigma_g^2 = \theta^2 + \gamma^2 - [\gamma \mathbb{E}(|Z_0|)]^2 + 2\,\theta\,\gamma\,\mathbb{E}(Z_0|Z_0|). \tag{2.7}$$

These properties are extremely important to prove the results stated in the sequel.

From now on, let $\lambda(\cdot)$ be the polynomial defined by

$$\lambda(z) = \frac{\alpha(z)}{\beta(z)}(1-z)^{-d} := \sum_{k=0}^{\infty} \lambda_{d,k} z^k, \quad \text{for all } |z| < 1, \tag{2.8}$$

where $\alpha(\cdot)$ and $\beta(\cdot)$ are defined in (2.3). Since it is assumed that $\beta(\cdot)$ has no roots in the closed disk $\{z : |z| \leq 1\}$, and also $\alpha(\cdot)$ and $\beta(\cdot)$ have no common roots, the function $\lambda(z)$ is analytic in the open disc $\{z : |z| < 1\}$ (if $d \leq 0$, in the closed disk $\{z : |z| \leq 1\}$). Therefore, it has a unique power series representation and the operator given in (2.5) can be rewritten as $\frac{\alpha(\mathcal{B})}{\beta(\mathcal{B})}(1-\mathcal{B})^{-d} = \sum_{k=0}^{\infty} \lambda_{d,k}\mathcal{B}^k = \lambda(\mathcal{B})$. This representation is more convenient and will be extensively used when characterizing the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$.

Theorem 2.1 bellow gives the convergence order of the coefficients $\lambda_{d,k}$, as k goes to infinity. This result plays an important role when choosing the truncation point in the series representation for simulation purposes.

Theorem 2.1. *Let $\lambda(\cdot)$ be the polynomial defined by (2.8). Then, for all $k \in \mathbb{N}$, the coefficients $\lambda_{d,k}$ satisfy*

$$\lambda_{d,k} \sim \frac{1}{\Gamma(d)\,k^{1-d}} \frac{\alpha(1)}{\beta(1)}, \quad \text{as } k \to \infty. \tag{2.9}$$

Consequently, $\lambda_{d,k} = O(k^{d-1})$, as k goes to infinity.

Proof: Denote $\beta(z)^{-1}$ by $f(z)$. Since $\beta(\cdot)$ has no roots in the closed disk $\{z : |z| \leq 1\}$, one has

$$\beta(z)^{-1} := f(z) = \sum_{k=0}^{\infty} f_k z^k, \quad \text{where } f_k = \frac{f^{(k)}(0)}{k!}, \text{ for all } k \in \mathbb{N}. \tag{2.10}$$

From expressions (2.2), (2.8) and (2.10) it follows that

$$\lambda(z) = \sum_{k=0}^{\infty} \Big[\sum_{i=0}^{\min\{p,k\}} (-\alpha_i) \Big(\sum_{j=0}^{k-i} \pi_{d,k-i-j} f_j \Big) \Big] z^k. \tag{2.11}$$

From (2.11), one has

$$\lambda_{d,k} = \sum_{i=0}^{\min\{p,k\}} (-\alpha_i) \Big(\sum_{j=0}^{k-i} \pi_{d,k-i-j} f_j \Big), \quad \text{for all } k \in \mathbb{N}.$$

In particular, $\lambda_{d,k} = \sum_{i=0}^{p} (-\alpha_i) \Big(\sum_{j=0}^{k-i} \pi_{d,j} f_{k-i-j} \Big)$, for all $k > p$.

Moreover, since $f_k \to 0$, as $k \to \infty$, it follows that for all $\varepsilon > 0$, there exists $k_0 > 0$, such that, for a given $m > 0$ and for all $k > k_0$, $|\pi_{d,j} f_{k-i-j}| < \frac{\varepsilon}{m}$, for all $0 \leq j \leq m$ and $0 \leq i \leq p$. Hence, for k sufficiently large,

$$\lambda_{d,k} \sim \sum_{i=0}^{p} (-\alpha_i) \Big(\sum_{j=m+1}^{k-i} \pi_{d,j} f_{k-i-j} \Big).$$

Notice that, since $\pi_{d,k} \sim \frac{1}{\Gamma(d)\,k^{1-d}}$, as $k \to \infty$, one can choose m_0 such that $\pi_{d,k} \sim \pi_{d,k-i} \sim \pi_{d,j}$, for all $m_0 < m+1 \leq j \leq k-i$ and $0 \leq i \leq p$. Consequently,

$$\lambda_{d,k} \sim \pi_{d,k} \Big(\sum_{i=0}^{p} (-\alpha_i) \Big) \Big(\sum_{j=0}^{k-i-(m+1)} f_j \Big) \sim \pi_{d,k} \Big(\sum_{i=0}^{p} (-\alpha_i) \Big) \Big(\sum_{j=0}^{\infty} f_j \Big).$$

However, $\sum_{j=0}^{\infty} f_j = f(1) = \dfrac{1}{\beta(1)}$ and $\pi_{d,k} \sim \frac{1}{\Gamma(d)\,k^{1-d}}$, as $k \to \infty$. So, we have

$$\lambda_{d,k} \sim \pi_{d,k}\,\frac{\alpha(1)}{\beta(1)} \sim \frac{1}{\Gamma(d)\,k^{1-d}}\,\frac{\alpha(1)}{\beta(1)}.$$

It follows that $\lambda_{d,k} \to 0$ and $\lambda_{d,k}k^{1-d} \to \frac{1}{\Gamma(d)}\frac{\alpha(1)}{\beta(1)}$, as $k \to \infty$. Hence, $\lambda_{d,k} = O(k^{d-1})$, as $k \to \infty$, which concludes the proof. ∎

Proposition 2.1 presents a recurrence formula for calculating the coefficients $\lambda_{d,k}$, for all $k \in \mathbb{N}$. This formula is used to generate the FIEGARCH time series in the simulation study presented in Section 4.

Proposition 2.1. *Let $\lambda(\cdot)$ be the polynomial defined by the expression (2.8). The coefficients $\lambda_{d,k}$, for all $k \in \mathbb{N}$, are given by*

$$\lambda_{d,0} = 1 \quad \textit{and} \quad \lambda_{d,k} = -\alpha_k^* + \sum_{i=0}^{k-1} \lambda_i \Big(\sum_{j=0}^{k-i} \beta_j^* \delta_{d,k-i-j} \Big), \ \textit{for all} \ \ k \geq 1, \tag{2.12}$$

where the coefficients $\delta_{d,k}$, for all $k \in \mathbb{N}$, are given in (2.1) and

$$\alpha_m^* := \begin{cases} \alpha_m, & \textit{if} \quad 0 \leq m \leq p; \\ 0, & \textit{if} \qquad m > p \end{cases} \qquad \textit{and} \quad \beta_m^* := \begin{cases} \beta_m, & \textit{if} \quad 0 \leq m \leq q; \\ 0, & \textit{if} \qquad m > q. \end{cases} \tag{2.13}$$

Proof: Let $\lambda(\cdot)$ be defined by (2.8). Consequently,

$$\alpha(z) = \beta(z)(1-z)^d \Big(\sum_{k=0}^{\infty} \lambda_{d,k} z^k \Big). \tag{2.14}$$

By defining β_k^* as in expression (2.13), for all $k \in \mathbb{N}$, and upon considering expression (2.1), observing that $\delta_{d,0} = -1 = \beta_0$, the right hand side of expression (2.14) can be rewritten as

$$\begin{aligned}\beta(z)(1-z)^d \Big(\sum_{k=0}^{\infty} \lambda_{d,k} z^k \Big) &= \Big[\sum_{k=0}^{\infty} \Big(\sum_{j=0}^{k} -\beta_j^* \delta_{d,k-j} \Big) z^k \Big] \Big(\sum_{k=0}^{\infty} \lambda_{d,k} z^k \Big) = \sum_{k=0}^{\infty} \Big[\sum_{i=0}^{k} \lambda_{d,i} \Big(-\sum_{j=0}^{k-i} \beta_j^* \delta_{d,k-i-j} \Big) \Big] z^k \\ &= \sum_{k=0}^{\infty} \Big[\lambda_{d,k} - \sum_{i=0}^{k-1} \lambda_{d,i} \Big(\sum_{j=0}^{k-i} \beta_j^* \delta_{d,k-i-j} \Big) \Big] z^k. \end{aligned} \tag{2.15}$$

Now, by setting α_k^* as in expression (2.13), for all $k \in \mathbb{N}$, from expression (2.15) one concludes that the equality (2.14) holds if and only if,

$$-\alpha_0^* = \lambda_{d,0} \quad \text{and} \quad -\alpha_k^* = \lambda_{d,k} - \sum_{i=0}^{k-1} \lambda_{d,i} \Big(\sum_{j=0}^{k-i} \delta_{d,k-i-j} \beta_j^* \Big), \quad \text{for all } k \geq 1.$$

Therefore, expression (2.12) holds. It is easy to see that by replacing the coefficients $\lambda_{d,k}$, given by (2.12), in the expression (2.15), for all $k \in \mathbb{N}$, we get $\sum_{k=0}^{\infty} (-\alpha_k^*) z^k = \alpha(z)$,which completes the proof. ∎

From Theorem 2.1 one concludes that $\{\lambda_{d,k}\}_{k \in \mathbb{N}} \in \ell^2$ if and only if $d < 0.5$ and $\{\lambda_{d,k}\}_{k \in \mathbb{N}} \in \ell^1$ whenever $d < 0$. Therefore, the random variable $\ln(\sigma_t^2)$, for all $t \in \mathbb{Z}$, is finite with probability one, if and only if $d < 0.5$. Moreover, since $\{g(Z_t)\}_{t \in \mathbb{Z}}$ is a white noise process, it follows immediately that $\{\ln(\sigma_t^2)\}_{t \in \mathbb{Z}}$ has an ARFIMA(q, d, p) representation. This fact is very useful for forecasting purposes (see Section 3). Also, the following properties hold

- if $d < 0.5$, the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is stationary (weakly and strictly) and ergodic;
- if $d \in (-1, 0.5)$ and $\alpha(z) \neq 0$, for $|z| \leq 1$, the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is invertible;
- if $d < 0.5$, the autocorrelation function of the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is such that

  $$\rho_{\ln(\sigma_t^2)}(h) \sim ch^{2d-1}, \quad \text{as} \;\; h \to \infty,$$

  where $c \neq 0$, and the spectral density function of the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is such that

  $$f_{\ln(\sigma_t^2)}(\lambda) = \frac{\sigma_g^2}{2\pi}\frac{|\alpha(e^{-i\lambda})|^2}{|\beta(e^{-i\lambda})|^2}|1-e^{-i\lambda}|^{-2d} \sim \frac{\sigma_g^2}{2\pi}\left(\frac{\alpha(1)}{\beta(1)}\right)^2\lambda^{-2d}, \quad \text{as } \lambda \to 0,$$

  where $\sigma_g^2 = \mathrm{Var}(g(Z_0))$ is given in (2.7);

Remark 2.3. In the literature one usually find that an ARFIMA(p, d, q) process is invertible for $|d| < 0.5$ (see, for instance, Brockwell and Davis, 1991). However, Bloomfield (1985) seems to be the first to notice that this range can be extended to $d \in (-1, 0.5)$, for an ARFIMA$(0, d, 0)$. Recently, Bondon and Palma (2007) show that this result actually holds for any ARFIMA(p, d, q).

As a consequence of these properties we have the following corollary that extends the results in Nelson (1991), where only the case $d = 0$ is considered.

Corollary 2.1. *Let* $\{X_t\}_{t\in\mathbb{Z}}$ *be a* FIEGARCH(p, d, q) *process, given in Definition 2.1. If* $d < 0.5$*, then* $\{X_t\}_{t\in\mathbb{Z}}$ *and* $\{\sigma_t^2\}_{t\in\mathbb{Z}}$ *are strictly stationary and ergodic processes.*

Proof: Suppose that $d < 0.5$. Then, the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is weakly stationary and the random variable $\ln(\sigma_t^2)$ is almost surely finite, for all $t \in \mathbb{Z}$. It follows that σ_t^2 and X_t are also almost surely finite, for all $t \in \mathbb{Z}$. Since $\sigma_t^2 = \exp\{\ln(\sigma_t^2)\}$, for all $t \in \mathbb{Z}$, and the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is strictly stationary and ergodic, it follows that $\{\sigma_t^2\}_{t\in\mathbb{Z}}$ is also a strictly stationary and ergodic process. From expressions (2.4), (2.5) and (2.8), one has

$$X_t = Z_t \exp\left\{\frac{1}{2}\Big(\omega + \sum_{k=0}^{\infty}\lambda_{d,k}g(Z_{t-1-k})\Big)\right\}, \text{ for all } t \in \mathbb{Z}.$$

Since $\{Z_t\}_{t\in\mathbb{Z}}$ is a sequence of i.i.d. random variables, it follows that $\{X_t\}_{t\in\mathbb{Z}}$ is a strictly stationary and ergodic process. ∎

Corollary 2.1 shows that $\{X_t\}_{t\in\mathbb{Z}}$ and $\{\sigma_t^2\}_{t\in\mathbb{Z}}$ are strictly stationary and ergodic processes. However, this does not imply weakly stationarity when the random variable Z_t, for $t \in \mathbb{Z}$, is such that either its mean or its variance is not finite. Theorem 2.2 in Nelson (1990) considers the generalized error distribution (GED) function and proves the existence of the moment of order $r > 0$, for the random variables X_t and σ_t^2, for all $t \in \mathbb{Z}$, when the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is defined in terms of a square summable sequence of coefficients. Theorem 2.2 bellow shows that, whenever $d < 0.5$, the theorem 2.2 in Nelson (1990) also holds for FIEGARCH(p, d, q) processes.

Theorem 2.2. *Let* $\{X_t\}_{t\in\mathbb{Z}}$ *be a* FIEGARCH(p, d, q) *process, given in Definition 2.1. Suppose* θ *and* γ *are not both equal to zero and* $d < 0.5$*. Also suppose that* $\{Z_t\}_{t\in\mathbb{Z}}$ *is a sequence of i.i.d. random variables with zero mean and variance equal to one, with* GED(v) *distribution function, where* $v > 1$*. Then, the processes* $\{X_t\}_{t\in\mathbb{Z}}$ *and* $\{\sigma_t^2\}_{t\in\mathbb{Z}}$ *are such that* $\mathbb{E}(X_t^r) < \infty$ *and* $\mathbb{E}([\sigma_t^2]^r) < \infty$*, for all* $t \in \mathbb{Z}$ *and* $r > 0$*.*

Proof: For any FIEGARCH(p,d,q) process, from expressions (2.5) and (2.8), $\ln(\sigma_t^2)$ can be rewritten as

$$\ln(\sigma_t^2) = \omega + \sum_{k=0}^{\infty} \lambda_{d,k} g(Z_{t-1-k}), \text{ for all } t \in \mathbb{Z}. \tag{2.16}$$

It is easy to see that the expression (2.16) is equivalent to expression (2.1) of Nelson (1991). Indeed, one just needs to set $\alpha_t = \omega$, for all $t \in \mathbb{Z}$, and $\beta_k = \lambda_{d,k-1}$, for all $k \geq 1$. Moreover, by definition, the polynomial $\beta(\cdot)$ has no roots in the closed disk $\{z : |z| \leq 1\}$. Therefore, if $d < 0.5$, then $\sum_{k=0}^{\infty} \lambda_{d,k}^2 < \infty$. Under these conditions, the result follows from Nelson (1991), theorem 2.2. ∎

Since $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA process, its properties are well known. However, in practice, this process cannot be directly observed and frequently, knowing its characteristics may not help to identify a model that fits the data. On the other side, by expression (2.4), one can write $\ln(X_t^2) = \ln(\sigma_t^2) + \ln(Z_t^2)$, for all $t \in \mathbb{Z}$. Since the process $\{X_t\}_{t\in\mathbb{Z}}$ is observable, $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is a known process.

Perez and Zaffaroni (2008) and Hurvich et al. (2005) consider models such that X_t can be written as in (2.4), but σ_t can have a more general definition than (2.5). While Perez and Zaffaroni (2008) consider maximum likelihood and Whittle's method of estimation in the class of exponential volatility models, especially the EGARCH ones, Hurvich et al. (2005) consider different semiparametric estimators of the memory parameter in general signal plus noise models. In both cases, to obtain an estimator by Whittle's method, the authors consider the spectral density function of $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$.

Here we consider only the case where X_t can be written as in (2.4), and σ_t is defined through the expression (2.5) and we present some properties of the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ such as stationarity and ergodicity. We also show that, under mild conditions, this process also has an ARFIMA representation. To the best of our knowledge, up to this date, no formal proves of these results are given in the literature of FIEGARCH(p,d,q) processes, specially the ARFIMA$(q,d,0)$ representation of $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$.

Theorem 2.3. *Let $\{X_t\}_{t\in\mathbb{Z}}$ be a FIEGARCH(p,d,q) process, given in Definition 2.1. If $d < 0.5$ and $\mathbb{E}([\ln(Z_t^2)]^2) < \infty$, for all $t \in \mathbb{Z}$, then the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is well defined and it is stationary (weakly and strictly) and ergodic. Moreover, the autocovariance function of $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is given by*

$$\gamma_{\ln(X^2)}(h) = \sigma_g^2 \sum_{k=0}^{\infty} \lambda_{d,k}\lambda_{d,k+|h|} + \mathrm{Var}(\ln(Z_t^2))\mathbb{I}_{\{0\}}(h) + \lambda_{d,|h|-1}\mathcal{K}\mathbb{I}_{\mathbb{Z}^*}(h), \quad \textit{for all } h \in \mathbb{Z}. \tag{2.17}$$

where σ_g^2 is given in (2.7) and $\mathcal{K} = \mathrm{Cov}\big(g(Z_0), \ln(Z_0^2)\big)$.

Proof: If $d < 0.5$, the random variable $\ln(\sigma_t^2)$ is finite with probability one, for all $t \in \mathbb{Z}$. If $\mathbb{E}([\ln(Z_0^2)]^2) < \infty$, one has $\mathbb{E}(|\ln(Z_0^2)|) < \infty$ and the random variable $|\ln(Z_t^2)|$ is also finite with probability one, for all $t \in \mathbb{Z}$. Which implies that the random variable $\ln(X_t^2)$ is finite with probability one, for all $t \in \mathbb{Z}$. Therefore, the stochastic process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is well defined. By definition, $\{Z_t\}_{t\in\mathbb{Z}}$ is a sequence of i.i.d. random variables and $\ln(\sigma_t^2)$ and $\ln(Z_t^2)$ are measurable functions of $\{Z_t\}_{t\in\mathbb{Z}}$. Therefore, the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is strictly stationary and ergodic (see Durrett, 1991).

Since $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is a strictly stationary process, to prove that it is also weakly stationary one only needs to show that $\mathbb{E}([\ln(X_t^2)]^2) < \infty$ or, equivalently, $\mathrm{Var}(\ln(X_t^2)) < \infty$, for all $t \in \mathbb{Z}$. Notice that, if $d < 0.5$, $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is weakly stationary which implies that $\mathrm{Var}(\ln(\sigma_t^2)) < \infty$ and, if $\mathbb{E}([\ln(Z_t^2)]^2) = \mathbb{E}([\ln(Z_0^2)]^2) < \infty$, then $\mathrm{Var}(\ln(Z_t^2)) < \infty$, for all $t \in \mathbb{Z}$. Moreover, from expressions (2.5) and (2.8), $\ln(\sigma_t^2)$ is a measurable function of $\{Z_s\}_{s\leq t-1}$. It follows that, $\ln(\sigma_t^2)$ and $\ln(Z_t^2)$ are independent, for all $t \in \mathbb{Z}$. Hence,

$$\mathrm{Var}(\ln(X_t^2)) = \mathrm{Var}(\ln(\sigma_t^2)) + \mathrm{Var}(\ln(Z_t^2)) < \infty, \quad \text{for all} \quad t \in \mathbb{Z}.$$

To complete the prove we need to show that the autocovariance function $\gamma_{\ln(X^2)}(h)$, for all $h \in \mathbb{Z}$, is given by expression (2.17). From the definition of $\ln(X_t^2)$, it follows that

$$\begin{aligned}\mathrm{Cov}\big(\ln(X_{t+h}^2), \ln(X_t^2)\big) &= \mathrm{Cov}\big(\ln(\sigma_{t+h}^2), \ln(\sigma_t^2)\big) + \mathrm{Cov}\big(\ln(Z_{t+h}^2), \ln(Z_t^2)\big) \\ &\quad + \mathrm{Cov}\big(\ln(\sigma_{t+h}^2), \ln(Z_t^2)\big) + \mathrm{Cov}\big(\ln(Z_{t+h}^2), \ln(\sigma_t^2)\big). \end{aligned} \tag{2.18}$$

Since $\{g(Z_t)\}_{t\in\mathbb{Z}}$ is a white noise process, from expression (2.16) one concludes that

$$\mathrm{Cov}\big(\ln(\sigma_{t+h}^2), \ln(\sigma_t^2)\big) = \sigma_g^2 \sum_{k=0}^{\infty} \lambda_{d,k}\lambda_{d,k+|h|}, \quad \text{for all } h \in \mathbb{Z}.$$

From the independence of the random variables $\ln(Z_t^2)$, for all $t \in \mathbb{Z}$, and from expression (2.16), we have

$$\mathrm{Cov}\big(\ln(Z_{t+h}^2), \ln(Z_t^2)\big) = \begin{cases} 0, & \text{if } h \neq 0; \\ \mathrm{Var}(\ln(Z_t^2)), & \text{if } h = 0 \end{cases} \quad \text{and} \quad \mathrm{Cov}\big(\ln(\sigma_{t+h}^2), \ln(Z_t^2)\big) = \begin{cases} \lambda_{d,h-1}\mathcal{K}, & \text{if } h > 0; \\ 0, & \text{if } h \leq 0. \end{cases}$$

where $\mathcal{K} = \mathrm{Cov}\big(g(Z_0), \ln(Z_0^2)\big)$. Since $\mathrm{Cov}\big(\ln(Z_{t+h}^2), \ln(\sigma_t^2)\big) = \mathrm{Cov}\big(\ln(\sigma_{u-h}^2), \ln(Z_u^2)\big)$, with $u = t + h$, one concludes that

$$\mathrm{Cov}\big(\ln(\sigma_{t+h}^2), \ln(Z_t^2)\big) + \mathrm{Cov}\big(\ln(Z_{t+h}^2), \ln(\sigma_t^2)\big) = \mathrm{Cov}\big(\ln(\sigma_{t+|h|}^2), \ln(Z_t^2)\big) = \lambda_{d,|h|-1}\mathcal{K}\mathbb{I}_{\mathbb{Z}^*}(h).$$

By replacing these results on expression (2.18) we conclude that the autocovariance function of $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is given by (2.17). ∎

Figure 2.2 (a) presents the theoretical autocovariance function of the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$, where $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH$(0, d, 1)$ process, with the same parameter values considered in Figures 2.1 and 2.3. Figure 2.2 (b) shows the sample autocovariance function of the time series $\{\ln(X_t^2)\}_{t=1}^{2000}$, where $\{X_t\}_{t=1}^{2000}$ is the simulated time series presented in Figure 2.1. Figure 2.2 (c) presents the sample autocovariance function of the time series $\{\ln(r_t^2)\}_{t=1}^{n}$, where $\{r_t\}_{t=1}^{n}$ is the Bovespa index log-returns time series (see Section 5). By comparing the three graphs in Figure 2.2, one concludes that all three functions present a similar behavior. Since the sample autocovariance function is an estimator of the theoretical autocovariance function, it is expected that their graphics will have the same behavior. The similarity between the decay in the graphs in Figure 2.2 (b) and (c) indicates that a FIEGARCH model seems appropriate for fitting the Bovespa index log-returns time series.

It is known that $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA(q, d, p) process. The following theorem shows that, under mild conditions, $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA$(q, d, 0)$ process with correlated innovations. In what follows, let $\phi(\cdot)$ be the polynomial defined by

$$\phi(z) := \beta(z)(1 - z)^d = \sum_{k=0}^{\infty} \phi_k z^k, \quad \text{for } |z| < 1. \tag{2.19}$$

Theorem 2.4. *Let $\{X_t\}_{t\in\mathbb{Z}}$ be a* FIEGARCH(p, d, q) *process, given in Definition 2.1. Suppose $|d| < 0.5$ and $\mathbb{E}([\ln(Z_0^2)]^2) < \infty$. Then, $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is an* ARFIMA$(q, d, 0)$ *process given by*

$$\beta(\mathcal{B})(1 - \mathcal{B})^d(\ln(X_t^2) - \omega) = \varepsilon_t, \quad \text{for all } t \in \mathbb{Z},$$



(a) $\gamma_{\ln(X^2)}(\cdot)$ (b) $\hat{\gamma}_{\ln(X^2)}(\cdot)$ (c) $\hat{\gamma}_{\ln(r^2)}(\cdot)$

Figure 2.2: (a) Theoretical autocovariance function of the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$, where $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH$(0,d,1)$ process and (b) sample autocovariance function of a time series $\{\ln(X_t^2)\}_{t=1}^{2000}$ derived from that FIEGARCH$(0,d,1)$ process; (c) sample autocovariance function of the time series $\{\ln(r_t^2)\}_{t=1}^{1717}$, where $\{r_t\}_{t=1}^{1717}$ is the Bovespa index log-returns time series.

where $\{\varepsilon_t\}_{t\in\mathbb{Z}}$ *is a stochastic process with zero mean and autocovariance function* $\gamma_\varepsilon(\cdot)$ *given by*

$$\gamma_\varepsilon(h) = \begin{cases} \sigma_g^2 \displaystyle\sum_{i=|h|}^{p} \alpha_i\alpha_{i-|h|} + \mathcal{K}\sum_{i=0}^{p}\alpha_i\phi_{i+|h|+1} + \mathcal{K}\sum_{i=|h|-1}^{p}\alpha_i\phi_{i-|h|+1} + \sigma_\ell^2\sum_{i=|h|}^{\infty}\phi_i\phi_{i-|h|}, & \text{if} \quad 0\leq |h| \leq p;\\ \mathcal{K}\alpha_p\phi_1 + \sigma_\ell^2 \displaystyle\sum_{i=p+1}^{\infty}\phi_i\phi_{i-(p+1)}, & \text{if} \quad |h| = p+1;\\ \sigma_\ell^2 \displaystyle\sum_{i=|h|}^{\infty}\phi_i\phi_{i-|h|}, & \text{if} \quad |h| > p+1, \end{cases} \tag{2.20}$$

with σ_g^2 *given in* (2.7), $\mathcal{K} = \mathrm{Cov}\big(g(Z_0), \ln(Z_0^2)\big)$, α_i *is given in* (2.3), *for all* $i \in \{0,\cdots,p\}$, ϕ_k *is defined in* (2.19), *for* $k\in\mathbb{N}$, *and* $\sigma_\ell^2 := \mathrm{Var}(\ln(Z_0^2))$.

Proof: Let $\{X_t\}_{t\in\mathbb{Z}}$ be a FIEGARCH process. From expressions (2.4) and (2.5) we have

$$\beta(\mathcal{B})(1-\mathcal{B})^d(\ln(X_t^2)-\omega) = \varepsilon_t, \quad \text{for all } t\in\mathbb{Z},$$

where

$$\varepsilon_t = \alpha(\mathcal{B})g(Z_{t-1}) + \beta(\mathcal{B})(1-\mathcal{B})^d\ln(Z_t^2), \text{ for all } t\in\mathbb{Z}. \tag{2.21}$$

In particular, if $d>0$, we have $\beta(\mathcal{B})(1-\mathcal{B})^d\omega = 0$ and $\beta(\mathcal{B})(1-\mathcal{B})^d\ln(X_t^2) = \varepsilon_t$, for all $t\in\mathbb{Z}$.

Now, suppose that $\mathbb{E}([\ln(Z_0^2)]^2) < \infty$. Since $\{Z_t\}_{t\in\mathbb{Z}}$ is a sequence of i.i.d. random variables and $0 \leq |\mathbb{E}(\ln(Z_0^2))| \leq \mathbb{E}(|\ln(Z_0^2)|) \leq [\mathbb{E}([\ln(Z_0^2)]^2)]^{1/2}$, one concludes that $\mathbb{E}(\ln(Z_t^2)) = \mathbb{E}(\ln(Z_0^2)) < \infty$, for all $t\in\mathbb{Z}$. Therefore, $\beta(\mathcal{B})(1-\mathcal{B})^d\mathbb{E}(\ln(Z_t^2)) = 0$ and $\alpha(\mathcal{B})\mathbb{E}(g(Z_{t-1})) = 0$. Consequently, $\mathbb{E}(\varepsilon_t) = 0$, for all $t\in\mathbb{Z}$.

Let $\phi(\cdot)$ be defined by expression (2.19). Assume, for the moment, that $\mathrm{Var}(\varepsilon_t^2) < \infty$, for all $t\in\mathbb{Z}$. It follows that

$$\begin{aligned} \mathrm{Cov}(\varepsilon_{t+h},\varepsilon_t) = {} & \mathrm{Cov}\big(\alpha(\mathcal{B})g(Z_{t+h-1}), \alpha(\mathcal{B})g(Z_{t-1})\big) + \mathrm{Cov}\big(\phi(\mathcal{B})\ln(Z_{t+h}^2), \phi(\mathcal{B})\ln(Z_t^2)\big)\\ & + \mathrm{Cov}\big(\alpha(\mathcal{B})g(Z_{t+h-1}), \phi(\mathcal{B})\ln(Z_t^2)\big) + \mathrm{Cov}\big(\phi(\mathcal{B})\ln(Z_{t+h}^2), \alpha(\mathcal{B})g(Z_{t-1})\big). \end{aligned} \tag{2.22}$$

Since $\{g(Z_t)\}_{t\in\mathbb{Z}}$ is a white noise process we have

$$\mathrm{Cov}\big(\alpha(\mathcal{B})g(Z_{t+h-1}), \alpha(\mathcal{B})g(Z_{t-1})\big) = \begin{cases} \mathrm{Var}(g(Z_0)) \displaystyle\sum_{i=|h|}^{p}\alpha_i\alpha_{i-|h|}, & \text{if} \quad |h|\leq p;\\ 0, & \text{if} \quad |h| > p, \end{cases}$$

which does not depend on $t \in \mathbb{Z}$. From the independence of the random variables Z_t, for all $t \in \mathbb{Z}$, one has

$$\operatorname{Cov}\big(\alpha(\mathcal{B})g(Z_{t+h-1}), \phi(\mathcal{B})\ln(Z_t^2)\big) = \begin{cases} \mathcal{K}\displaystyle\sum_{i=0}^{p} \alpha_i\phi_{i-h+1}, & \text{if} \quad h < 1; \\ \mathcal{K}\displaystyle\sum_{i=h-1}^{p} \alpha_i\phi_{i-h+1}, & \text{if} \quad 1 \leq h \leq p+1; \\ 0, & \text{if} \quad h > p+1 \end{cases}$$

and

$$\operatorname{Cov}\big(\phi(\mathcal{B})\ln(Z_{t+h}^2), \alpha(\mathcal{B})g(Z_{t-1})\big) = \begin{cases} 0, & \text{if} \quad h < -(p+1); \\ \mathcal{K}\displaystyle\sum_{i=|h|-1}^{p} \alpha_i\phi_{i+h+1}, & \text{if} \quad -(p+1) \leq h \leq -1; \\ \mathcal{K}\displaystyle\sum_{i=0}^{p} \alpha_i\phi_{i+h+1}, & \text{if} \quad h > -1, \end{cases}$$

where $\mathcal{K} := \operatorname{Cov}\big(g(Z_0), \ln(Z_0^2)\big)$ does not depend on $t \in \mathbb{Z}$. Also, from the independence of the random variables $\ln(Z_t^2)$, for all $t \in \mathbb{Z}$, we have

$$\operatorname{Cov}\big(\phi(\mathcal{B})\ln(Z_{t+h}^2), \phi(\mathcal{B})\ln(Z_t^2)\big) = \operatorname{Var}(\ln(Z_0^2)) \sum_{i=|h|}^{\infty} \phi_i\phi_{i-|h|}, \quad \text{for all } h \in \mathbb{Z},$$

which does not depend on $t \in \mathbb{Z}$.

Therefore, all four terms in expression (2.22) do not depend on $t \in \mathbb{Z}$ and expression (2.20) holds. Now, to validate expression (2.20) we only need to show that $\operatorname{Var}(\varepsilon_t) < \infty$, for all $t \in \mathbb{Z}$. Notice that, since $\mathbb{E}(\varepsilon_t) = 0$, it follows that $\mathbb{E}(\varepsilon_t^2) = \operatorname{Var}(\varepsilon_t) = \gamma_\varepsilon(0)$. Upon replacing $h = 0$ in (2.22) one obtains

$$\gamma_\varepsilon(0) = \operatorname{Var}(g(Z_0)) \sum_{i=0}^{p} \alpha_i^2 + 2\mathcal{K}\sum_{i=0}^{p} \alpha_i\phi_{i+1} + \operatorname{Var}(\ln(Z_0^2)) \sum_{i=0}^{\infty} \phi_i^2.$$

By hypothesis, $\mathbb{E}([\ln(Z_0^2)]^2) < \infty$ and $d \in (-0.5, 0.5)$. It follows that $\operatorname{Var}(\ln(Z_0^2)) < \infty$ and $\sum_{i=0}^{\infty} \phi_i^2 < \infty$. We also know that $\operatorname{Var}(g(Z_0)) < \infty$. In order to show that $\mathcal{K} < \infty$, notice that $\mathcal{K} := \operatorname{Cov}\big(g(Z_0), \ln(Z_0^2)\big) = \mathbb{E}\big(g(Z_0)\ln(Z_0^2)\big)$ and, since $\mathbb{E}(Z_0^2) = 1$ and $\operatorname{Var}(\ln(Z_0^2)) < \infty$, from Hölder's inequality, we have $\mathbb{E}(|Z_0|) < \infty$ and $\mathbb{E}(\ln(Z_0^2)) < \infty$. Then, from (2.6) it follows that

$$\mathbb{E}\big(g(Z_0)\ln(Z_0^2)\big) = \theta\, \mathbb{E}\big(Z_0 \ln(Z_0^2)\big) + \gamma \mathbb{E}\big(|Z_0| \ln(Z_0^2)\big) - c,$$

where $c := \gamma\, \mathbb{E}(|Z_0|)\mathbb{E}(\ln(Z_0^2)) < \infty$. By using the fact that $2ab \leq a^2 + b^2$, for all $a, b \in \mathbb{R}$, one concludes that

$$\big|\mathbb{E}\big(Z_t \ln(Z_t^2)\big)\big| \leq \frac{1}{2}\big[\mathbb{E}(Z_t^2) + \mathbb{E}\big(\ln(Z_t^2)\big)\big] < \infty \quad \text{and} \quad \big|\mathbb{E}\big(|Z_t| \ln(Z_t^2)\big)\big| \leq \frac{1}{2}\big[\mathbb{E}(Z_t^2) + \mathbb{E}\big(\ln(Z_t^2)\big)\big] < \infty.$$

Hence $\mathbb{E}\big(g(Z_0)\ln(Z_0^2)\big) < \infty$ and, consequently, $\operatorname{Cov}\big(g(Z_0), \ln(Z_0^2)\big) < \infty$ and $\gamma_\varepsilon(0) < \infty$. Therefore, the result follows. ∎

Remark 2.4. From Theorem 2.4 it follows that all classical results related to the stationarity and ergodicity of an ARFIMA process (see Lopes, 2008 for ARFIMA processes with non-correlated innovations) can be extended to the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$, where $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH(p, d, q) process.

The spectral density function of the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is given by (see Hurvich et al., 2005)

$$f_{\ln(X_t^2)}(\lambda) = f_{\ln(\sigma_t^2)}(\lambda) + \frac{\mathcal{K}}{\pi}\Re\big(e^{-\mathrm{i}\lambda}\Lambda(\lambda)\big) + f_{\ln(Z_t^2)}(\lambda), \quad \text{for all } \lambda \in [0, \pi], \tag{2.23}$$

where $\mathcal{K} = \mathrm{Cov}\big(g(Z_0), \ln(Z_0^2)\big)$, $\Re(z)$ is the real part of z and $\Lambda(z) := \lambda(e^{-\mathrm{i}z})$.

As an illustration, Figure 2.3 (a) shows the spectral density function of the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$, where $\{X_t\}_{t\in\mathbb{Z}}$ is any FIEGARCH$(0,d,1)$ process with $d = 0.3578$, $\theta = -0.1661$, $\gamma = 0.2792$, $\omega = -7.2247$ and $\beta_1 = 0.6860$. Figure 2.3 (b) shows the periodogram function of the time series $\{\ln(X_t^2)\}_{t=1}^{2000}$, where $\{X_t\}_{t=1}^{2000}$ is the simulated time series presented in Figure 2.1 (a). Figure 2.3 (c) shows the periodogram function of the time series $\{\ln(r_t^2)\}_{t=1}^{1717}$, where $\{r_t\}_{t=1}^{1717}$ is the Bovespa index log-returns time series. One observes that the graphs in Figure 2.3 (b) and (c) present similar behavior, indicating that a FIEGARCH model may be adequate to fit the data.



(a) $f_{\ln(X_t^2)}(\cdot)$ (b) $I_{\ln(X_t^2)}(\cdot)$ (c) $I_{\ln(r_t^2)}(\cdot)$

Figure 2.3: (a) Theoretical spectral density function of the process $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$, where $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH$(0,d,1)$ process with $d = 0.3578$, $\theta = -0.1661$, $\gamma = 0.2792$, $\omega = -7.2247$ and $\beta_1 = 0.6860$, (b) periodogram function related to a time series $\{X_t\}_{t=1}^{2000}$ derived from this FIEGARCH$(0,d,1)$ process; (c) the periodogram function related to the time series $\{\ln(r_t^2)\}_{t=1}^{1717}$, where $\{r_t\}_{t=1}^{1717}$ is the Bovespa index log-returns time series.

The following proposition presents the expression of the kurtosis and the asymmetry measures for any stationary FIEGARCH(p,d,q) process. This result generalizes the one in Ruiz and Veiga (2008) where the authors consider only the particular case $p = 0$ and $q = 1$.

Proposition 2.2. *Let* $\{X_t\}_{t\in\mathbb{Z}}$ *be a stationary* FIEGARCH(p,d,q) *process, given in Definition 2.1. If* $\mathbb{E}(Z_0^3) < \infty$, *the asymmetry measure of* $\{X_t\}_{t\in\mathbb{Z}}$ *is given by*

$$A_X = \mathbb{E}(Z_0^3)\,\frac{\displaystyle\prod_{k=0}^{\infty}\mathbb{E}\Big(\exp\Big\{\frac{3}{2}\lambda_{d,k}\,g(Z_0)\Big\}\Big)}{\Big[\displaystyle\prod_{k=0}^{\infty}\mathbb{E}\big(\exp\{\lambda_{d,k}\,g(Z_0)\}\big)\Big]^{3/2}}$$

and, if $\mathbb{E}(Z_0^4) < \infty$, *the kurtosis measure of* $\{X_t\}_{t\in\mathbb{Z}}$ *is given by*

$$K_X = \mathbb{E}(Z_0^4)\,\frac{\displaystyle\prod_{k=0}^{\infty}\mathbb{E}\big(\exp\{2\lambda_{d,k}\,g(Z_0)\}\big)}{\Big[\displaystyle\prod_{k=0}^{\infty}\mathbb{E}\big(\exp\{\lambda_{d,k}\,g(Z_0)\}\big)\Big]^{2}},$$

where $\lambda_{d,k}$ *are given in* (2.8) *and* $g(\cdot)$ *is defined by* (2.6).

Proof: Let $\{X_t\}_{t\in\mathbb{Z}}$ be any stationary FIEGARCH(p,d,q) process and $\lambda(\cdot)$ be the polynomial defined by (2.8). Since, by definition, σ_t and Z_t are independent, for all $t \in \mathbb{Z}$, it follows that

$$\mathbb{E}(X_t^r) = \mathbb{E}(X_0^r) = \mathbb{E}(\sigma_0^r)\mathbb{E}(Z_0^r), \quad \text{for all } t \in \mathbb{Z} \text{ and } r > 0.$$

In particular $\mathbb{E}(Z_t^2) = \mathbb{E}(Z_0^2) = 1$ and $\mathbb{E}(X_t^2) = \mathbb{E}(X_0^2) = \mathbb{E}(\sigma_0^2 Z_0^2) = \mathbb{E}(\sigma_0^2)$.

Since $\{g(Z_t)\}_{t\in\mathbb{Z}}$ is a sequence of i.i.d. random variables, from (2.5) it follows that

$$\mathbb{E}(\sigma_0^r) = e^{\frac{r\omega}{2}} \prod_{k=0}^{\infty} \mathbb{E}\Big(\exp\Big\{ \frac{r}{2}\lambda_{d,k} g(Z_0) \Big\} \Big), \quad \text{for all } r > 0. \tag{2.24}$$

Also, it is known that $\mathbb{E}(X_0) = 0$ and $\mathbb{E}(Z_0^2) = 1$, for all $t \in \mathbb{Z}$. Therefore, the asymmetry and the kurtosis measures of $\{X_t\}_{t\in\mathbb{Z}}$ are given, respectively, by

$$A_X = \frac{\mathbb{E}(X_t^3)}{[\mathbb{E}(X_t^2)]^{3/2}} = \frac{\mathbb{E}(\sigma_t^3)\mathbb{E}(Z_t^3)}{[\mathbb{E}(\sigma_t^2)]^{3/2}} \quad \text{and} \quad K_X = \frac{\mathbb{E}(X_t^4)}{[\mathbb{E}(X_t^2)]^2} = \frac{\mathbb{E}(\sigma_t^4)\mathbb{E}(Z_t^4)}{[\mathbb{E}(\sigma_t^2)]^2}. \tag{2.25}$$

Upon replacing (2.24) in (2.25) we concludes the proof. ∎

Figure 2.4 shows the theoretical value of the kurtosis measure, as a function of the parameter d, for any FIEGARCH(0,d,1) process, with Gaussian noise and parameters $\theta = -0.1661$, $\gamma = 0.2792$, $\omega = -7.2247$ and (a) $\beta_1 = 0.6860$ (b) $\beta_1 = -0.6860$. The parameter values considered in Figure 2.4 (a) are the same ones (except for d) considered in Figure 2.1 (a). For the specific model considered in Figure 2.1, $d = 0.3578$ and the theoretical value of the kurtosis measure is 5.6733. The sample kurtosis value of the simulated time series presented in Figure 2.1 (a) is 5.3197, which is very close to the theoretical one. It is easy to see that, while in Figure 2.4 (a) the kurtosis values increase exponentially as d increases, in Figure 2.4 (b) the kurtosis values decrease for $-0.5 \leq d < 0.3$ and increase for $0.3 \leq d \leq 0.5$.



(a) $\beta_1 = 0.6860$ (b) $\beta_1 = -0.6860$

Figure 2.4: (a) Kurtosis measure of a FIEGARCH$(0, d, 1)$ process as a function of the parameter d with $\theta = -0.1661$, $\gamma = 0.2792$, $\omega = -7.2247$ and $\beta_1 = 0.6860$; (b) Kurtosis measure of a FIEGARCH$(0, d, 1)$ process with the same parameters as in (a) but with β_1 replaced by $\beta_1 = -0.6860$.

3 Forecasting

Let $\{X_t\}_{t\in\mathbb{Z}}$ be a FIEGARCH(p, d, q) process, given in Definition 2.1, and $\{X_t\}_{t=1}^n$ a time series obtained from this process. In this section, we prove that $\{X_t\}_{t\in\mathbb{Z}}$ is a martingale difference with respect to the filtration $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$, where $\mathcal{F}_t := \sigma(\{Z_s\}_{s\leq t})$, and we provide the h-step ahead forecast for the process $\{X_t\}_{t\in\mathbb{Z}}$. Since the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$, defined by (2.5), has an ARFIMA(q, d, p) representation, the h-step ahead forecasting for this process and its mean square error value can be easily obtained (for instance, see Lopes, 2008 and Reisen and Lopes, 1999). This fact is used to provide an h-step ahead forecast for $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ and the mean square error of forecasting. We also consider the fact that $\mathbb{E}(X_t^2) = \mathbb{E}(\sigma_t^2)$, for all $t \in \mathbb{Z}$, to provide an h-step ahead forecast for both processes, $\{X_t^2\}_{t\in\mathbb{Z}}$ and $\{\sigma_t^2\}_{t\in\mathbb{Z}}$, based on the predictions obtained from the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$. The notation used in this section is introduced bellow.

Remark 3.1. Let Y_t, for $t \in \mathbb{Z}$, denote any random variable defined here. In the sequel we consider the following notation:

- we use the symbol "$\hat{\ }$" to denote the h-ahead step forecast defined in terms of the conditional expectation, that is, $\hat{Y}_{t+h} = \mathbb{E}(Y_{t+h}|\mathcal{F}_t)$. Notice that this is the best linear predictor in terms of mean square error value. The symbols "$\tilde{\ }$" and "$\check{\ }$" are used to denote alternative estimators (e.g. $\tilde{Y}_{t+h}$ and $\check{Y}_{t+h}$);
- for simplicity of notation, for the h-step ahead forecast of $\ln(Y_{t+h})$, we write $\hat{\ln}(Y_{t+h})$ instead of $\widehat{\ln(Y_{t+h})}$ (analogously for "$\tilde{\ }$" and "$\check{\ }$");
- we follow the approach usually considered in the literature and denote the h-ahead step forecast Y_{t+h}^2 as $\hat{Y}_{t+h}^2$ instead of $\widehat{Y_{t+h}^2}$. If necessary, to avoid confusion, we will denote the square of $\hat{Y}_{t+h}$ as $(\hat{Y}_{t+h})^2$ (analogously for "$\tilde{\ }$" and "$\check{\ }$").

The following lemma shows that a FIEGARCH(p,d,q) process is a martingale difference with respect to $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$. This result is useful in the proof of Lemma 3.2 that presents the h-step ahead forecast of X_{n+h}, for a fixed value of $n \in \mathbb{Z}$ and all $h \geq 1$, and the 1-step ahead forecast of X_{n+1}^2, given $\mathcal{F}_n$.

Lemma 3.1. *Let $\{X_t\}_{t\in\mathbb{Z}}$ be a* FIEGARCH(p,d,q) *process, given in Definition 2.1 and $\mathcal{F}_t := \sigma(\{Z_s\}_{s\leq t})$. Then, the process $\{X_t\}_{t\in\mathbb{Z}}$ is a martingale difference with respect to $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$.*

Proof: From definition, σ_t is a $\mathcal{F}_{t-1}$-measurable function. Moreover, $\mathbb{E}(X_t) = \mathbb{E}(\mathbb{E}(X_t|\mathcal{F}_{t-1}))$ and $\mathbb{E}(X_t|\mathcal{F}_{t-1}) = \mathbb{E}(\sigma_t Z_t|\mathcal{F}_{t-1}) = \sigma_t \mathbb{E}(Z_t|\mathcal{F}_{t-1}) = 0$. Therefore, the process $\{X_t\}_{t\in\mathbb{Z}}$ is a martingale difference with respect to $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$. ∎

Lemma 3.2. *Let $\{X_t\}_{t\in\mathbb{Z}}$ be a stationary* FIEGARCH(p,d,q) *process, given by Definition 2.1. Then, for any fixed $n \in \mathbb{Z}$, the h-step ahead forecast of X_{n+h}, for all $h > 0$ and the* 1*-step ahead forecast of X_{n+1}^2, given $\mathcal{F}_n$, are, respectively, $\hat{X}_{n+h} = 0$ and $\hat{X}_{n+1}^2 = \sigma_{n+1}^2$.*

Proof: From Lemma 3.1, a FIEGARCH(p,d,q) process is a martingale difference. It follows that $\hat{X}_{n+h} = \mathbb{E}(X_{n+h}|\mathcal{F}_n) = 0$, for all $h > 0$. From definition, $\mathbb{E}(X_{n+1}^2|\mathcal{F}_n) = \sigma_{n+1}^2$. Therefore, the 1-step ahead forecast of X_{n+1}^2, given $\mathcal{F}_n$, is σ_{n+1}^2. Moreover, if $\mathbb{E}(X_t^4) < \infty$, for all $t \in \mathbb{Z}$, then, this is the best forecast value in mean square error sense. ∎

To obtain the h-step ahead forecast for X_{n+h}^2, notice that σ_t and Z_t are independent and so are σ_t^2 and Z_t^2, for all $t \in \mathbb{Z}$. Moreover, $\mathbb{E}(Z_{n+h}^2|\mathcal{F}_n) = \mathbb{E}(Z_{n+h}^2) = 1$, for all $h > 0$. It follows that

$$\hat{X}_{n+h}^2 := \mathbb{E}(X_{n+h}^2|\mathcal{F}_n) = \mathbb{E}(\sigma_{n+h}^2|\mathcal{F}_n) := \hat{\sigma}_{n+h}^2, \quad \text{for all} \quad h > 0.$$

While for ARCH/GARCH models, $\mathbb{E}(\sigma_{n+h}^2|\mathcal{F}_t)$ can be easily calculated, for FIEGARCH processes, what is easy to derive is the expression for the h-step ahead forecast for the process $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$, for any $h > 1$. The expressions for $\hat{\ln}(\sigma_{n+h}^2) := \mathbb{E}(\ln(\sigma_{n+h}^2)|\mathcal{F}_t)$ and for the mean square error of forecast are given in Proposition 3.1. We shall use this result to discuss the properties of the predictor obtained by considering $\check{\sigma}_{n+h}^2 := \exp\{\hat{\ln}(\sigma_{n+h}^2)\}$, for all $h > 0$.

Proposition 3.1. *Let $\{X_t\}_{t\in\mathbb{Z}}$ be a* FIEGARCH(p,d,q) *process, given by Definition 2.1. Then, the h-step ahead forecast $\hat{\ln}(\sigma_{n+h}^2)$ of $\ln(\sigma_{n+h}^2)$, given $\mathcal{F}_n = \sigma(\{Z_t\}_{t\leq n})$, $n \in \mathbb{N}$, is given by*

$$\hat{\ln}(\sigma_{n+h}^2) = \omega + \sum_{k=0}^{\infty} \lambda_{d,k+h-1}\, g(Z_{n-k}), \text{ for all } h > 0. \tag{3.1}$$

Moreover, the mean square error forecast is equal to zero, if $h = 1$*, and it is given by*

$$\mathbb{E}\big(\big[\ln(\sigma_{n+h}^2) - \hat{\ln}(\sigma_{n+h}^2)\big]^2\big) = \sigma_g^2 \sum_{k=0}^{h-2} \lambda_{d,k}^2, \quad \textit{if } h \geq 2, \tag{3.2}$$

where $\sigma_g^2 := \mathbb{E}([g(Z_0)]^2)$ *is given in* (2.7).

Proof: Let $\hat{\ln}(\sigma_{n+h}^2) := \mathbb{E}(\ln(\sigma_{n+h}^2)|\mathcal{F}_n)$. Note that $\mathbb{E}(g(Z_t)|\mathcal{F}_n) = \mathbb{E}(g(Z_t)) = 0$, for all $t > n$, and $\mathbb{E}(g(Z_t)|\mathcal{F}_n) = g(Z_t)$, for all $t \leq n$. By the expression (2.16), one has

$$\hat{\ln}(\sigma_{n+h}^2) = \omega + \sum_{k=0}^{\infty} \lambda_{d,k}\, \mathbb{E}(g(Z_{n+h-1-k})|\mathcal{F}_n) = \omega \; + \sum_{k=h-1}^{\infty} \lambda_{d,k}\, g(Z_{n+h-1-k}),$$

and expression (3.1) follows.

Since $\ln(\sigma_{n+h}^2)$ is a function of $\{g(Z_s)\}_{s \leq n+h-1}$ and $\{g(Z_t)\}_{t\in\mathbb{Z}}$ is a sequence of i.i.d. random variables with zero mean and variance $\sigma_g^2 := \mathbb{E}([g(Z_0)]^2)$, we conclude that

$$\mathbb{E}\big(\big[\ln(\sigma_{n+h}^2) - \hat{\ln}(\sigma_{n+h}^2)\big]^2\big) = \mathbb{E}\Big(\Big[\sum_{k=0}^{h-2} \lambda_{d,k}\, g(Z_{n+h-1-k})\Big]^2\Big) = \sigma_g^2 \sum_{k=0}^{h-2} \lambda_{d,k}^2, \quad \text{if } h \geq 2,$$

and zero if $h = 1$. ∎

In practice, $\mathbb{E}(\sigma_{n+h}^2|\mathcal{F}_t)$ cannot be easily calculated for FIEGARCH models and thus, a common approach is to predict σ_{n+h}^2 through the relation $\check{\sigma}_{n+h}^2 := \exp\{\hat{\ln}(\sigma_{n+h}^2)\}$, with $\hat{\ln}(\sigma_{n+h}^2)$ defined by (3.1), for all $h > 0$. As a consequence, a h-step ahead forecast for X_{n+h}^2 is defined as $\check{X}_{n+h}^2 := \check{\sigma}_{n+h}^2$ and a naive estimator for $\ln(X_{n+h}^2)$ is obtained by letting

$$\check{\ln}(X_{n+h}^2) := \ln(\check{X}_{n+h}^2) = \ln(\check{\sigma}_{n+h}^2) = \hat{\ln}(\sigma_{n+h}^2), \qquad \text{for all} \quad h > 0. \tag{3.3}$$

From expressions (2.4) and (3.3), it is obvious that $\check{\ln}(X_{n+h}^2)$ is a biased estimator for $\ln(X_{n+h}^2)$, whenever $\mathbb{E}(\ln(Z_{n+h}^2)) \neq 0$. Proposition 3.2 gives the mean square error forecast for the h-step ahead forecast of $\ln(X_{n+h}^2)$, defined through expression (3.3).

Proposition 3.2. *Let* $\check{\ln}(X_{n+h}^2)$*, for all* $h > 0$*, be the* h*-step ahead forecast of* $\ln(X_{n+h}^2)$*, given the filtration* $\mathcal{F}_n = \sigma(\{Z_s\}_{s\leq n})$*, defined by expression* (3.3)*. Then, the mean square error forecast is given by*

$$\mathbb{E}\big(\big[\ln(X_{n+h}^2) - \check{\ln}(X_{n+h}^2)\big]^2\big) = \sigma_g^2 \sum_{k=0}^{h-2} \lambda_{d,k}^2 + \mathbb{E}\big(\big[\ln(Z_{n+h}^2)\big]^2\big), \quad \textit{where } \sigma_g^2 := \mathbb{E}([g(Z_0)]^2).$$

Proof: By expression (3.3), $\check{\ln}(X_{n+h}^2) := \hat{\ln}(\sigma_{n+h}^2)$, for all $h > 0$. Thus, from expression (2.4) and from Proposition 3.1, we have

$$\begin{aligned}\mathbb{E}\big(\big[\ln(X_{n+h}^2) - \check{\ln}(X_{n+h}^2)\big]^2\big) &= \mathbb{E}\big(\big[\ln(X_{n+h}^2) - \ln(\hat{\sigma}_{n+h}^2)\big]^2\big) = \mathbb{E}\big(\big[\ln(\sigma_{n+h}^2) + \ln(Z_{n+h}^2) - \ln(\hat{\sigma}_{n+h}^2)\big]^2\big) \\ &= \mathbb{E}\Big(\Big[\sum_{k=0}^{h-2} \lambda_{d,k}\, g(Z_{n+h-1-k}) + \ln(Z_{n+h}^2)\Big]^2\Big). \end{aligned} \tag{3.4}$$

By expanding the right hand side of expression (3.4) and using the fact that $\{Z_t\}_{t\in\mathbb{Z}}$ is a sequence of i.i.d. random variables, the proposition follows immediately. ∎

Remark 3.2. If the values of X_t and σ_t are known only for $t \in \{1, \cdots, n\}$ then, the h-step ahead forecast $\hat{\ln}(\hat{\sigma}_{n+h}^2)$ of $\ln(\sigma_{n+h}^2)$, is approximated by

$$\hat{\ln}(\sigma^2_{n+h}) \simeq \omega + \sum_{k=0}^{n-1} \lambda_{d,k+h-1}\, g(Z_{n-k}), \quad \text{for all } h > 0,$$

and, by definition, the same approximation follows for $\check{\ln}(X^2_{n+h})$. It is easy to see that, in this case, the mean square error of forecast values for the processes $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ are given, respectively, by

$$\mathbb{E}\big(\big[\ln(\sigma^2_{n+h}) - \hat{\ln}(\sigma^2_{n+h})\big]^2\big) = \sigma_g^2\Big(\sum_{k=0}^{h-2}\lambda^2_{d,k} \;+ \sum_{k=n+h-1}^{\infty}\lambda^2_{d,k}\Big), \quad \text{and}$$

$$\mathbb{E}\big(\big[\ln(X^2_{n+h}) - \check{\ln}(X^2_{n+h})\big]^2\big) = \sigma_g^2\Big(\sum_{k=0}^{h-2}\lambda^2_{d,k} \;+ \sum_{k=n+h-1}^{\infty}\lambda^2_{d,k}\Big) + \mathbb{E}\big(\big[\ln(Z^2_{n+h})\big]^2\big), \quad \text{for all } h > 0.$$

From Jensen's inequality, one concludes that

$$\check{\sigma}^2_{n+h} := \exp\{\hat{\ln}(\sigma^2_{n+h})\} = \exp\{\mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_t\big)\} \le \mathbb{E}(\sigma^2_{n+h}|\mathcal{F}_n) := \hat{\sigma}^2_{n+h}, \quad \text{for all } h > 0,$$

so that $\mathbb{E}(\check{\sigma}^2_{n+h} - \sigma^2_{n+h}) = \mathbb{E}(\mathbb{E}(\check{\sigma}^2_{n+h} - \sigma^2_{t+h}|\mathcal{F}_n)) = \mathbb{E}(\check{\sigma}^2_{n+h} - \hat{\sigma}^2_{n+h}) \le 0$, for all $h > 0$. In fact, from (2.9) and (3.1), we have

$$\check{\sigma}^2_{n+h} := \exp\{\hat{\ln}(\sigma^2_{n+h})\} = \exp\Big\{\omega + \sum_{k=0}^{\infty}\lambda_{d,k+h-1}\, g(Z_{n-k})\Big\} \xrightarrow{h\to\infty} e^{\omega} = \exp\{\mathbb{E}\big(\ln(\sigma_0^2)\big)\}. \tag{3.5}$$

Another h-step ahead predictor for σ^2_{n+h} can be defined as follows. Consider an order 2 Taylor's expansion of the exponential function and write

$$\begin{aligned}\sigma^2_{n+h} &= \exp\big\{\mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big\} + \big[\ln(\sigma^2_{n+h}) - \mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big]\exp\big\{\mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big\}\\ &\quad + \frac{1}{2}\big[\ln(\sigma^2_{n+h}) - \mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big]^2\exp\big\{\mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big\} + R_{n+h}, \quad \text{for all } h > 0,\end{aligned} \tag{3.6}$$

From expression (3.6), a natural choice is to define a h-step ahead predictor for σ^2_{n+h} as

$$\tilde{\sigma}^2_{n+h} := \exp\big\{\mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big\} + \frac{1}{2}\mathbb{E}\big(\big[\ln(\sigma^2_{n+h}) - \mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big]^2\big)\exp\big\{\mathbb{E}\big(\ln(\sigma^2_{n+h})|\mathcal{F}_n\big)\big\}, \tag{3.7}$$

for all $h > 0$.

From expressions (3.1) and (3.2) one concludes that $\check{\sigma}^2_{n+h}$ and $\tilde{\sigma}^2_{n+h}$ are related through the equation

$$\tilde{\sigma}^2_{n+h} = \begin{cases} \exp\{\hat{\ln}(\sigma^2_{n+h})\} = \check{\sigma}^2_{n+h}, & \text{if} \quad h = 1;\\[2ex] \exp\{\hat{\ln}(\sigma^2_{n+1})\}\Big(1 + \dfrac{1}{2}\sigma_g^2\displaystyle\sum_{k=0}^{h-2}\lambda^2_{d,k}\Big) = \check{\sigma}^2_{n+h}\Big(1 + \dfrac{1}{2}\sigma_g^2\displaystyle\sum_{k=0}^{h-2}\lambda^2_{d,k}\Big), & \text{if} \quad h > 1.\end{cases} \tag{3.8}$$

Since σ_{t+1} is a $\mathcal{F}_t$-measurable random variable, for all $t \in \mathbb{Z}$, we have $\mathbb{E}(\tilde{\sigma}^2_{n+1} - \sigma^2_{n+1}) = \mathbb{E}(\check{\sigma}^2_{n+1} - \sigma^2_{n+1}) = 0$. From equation (3.6), we easily conclude that, for all $h > 1$,

$$\mathbb{E}(\tilde{\sigma}^2_{n+h} - \sigma^2_{n+h}) = -\mathbb{E}(R_{n+h}) \quad \text{and} \quad \mathbb{E}(\check{\sigma}^2_{n+h} - \sigma^2_{n+h}) = -\Big(1 + \frac{1}{2}\sigma_g^2\sum_{k=0}^{h-2}\lambda^2_{d,k}\Big)\mathbb{E}(\check{\sigma}^2_{n+h}) - \mathbb{E}(R_{n+h}).$$

Therefore, the relation between the bias for the estimators $\check{\sigma}^2_{n+h}$ and $\tilde{\sigma}^2_{n+h}$ is given by

$$\mathbb{E}(\tilde{\sigma}^2_{n+h} - \sigma^2_{n+h}) = \mathbb{E}(\check{\sigma}^2_{n+h} - \sigma^2_{n+h}) + \mathbb{E}(\check{\sigma}^2_{n+h})\Big(1 + \frac{1}{2}\sigma_g^2\sum_{k=0}^{h-2}\lambda^2_{d,k}\Big), \quad \text{for all } h > 1.$$

In Section 4 we analyze the performance of $\tilde{\sigma}^2_{n+h}$ through a Monte Carlo simulation study.

4 Simulation Study

In this section we present a Monte Carlo simulation study to analyze the performance of quasi-likelihood estimator and also the forecasting on FIEGARCH(p, d, q) processes. The set of parameters considered in this study is the same as in Prass and Lopes (2011a) and Prass and Lopes (2011b), except for model M5 (see Table 4.1). Prass and Lopes (2011a) present a Monte Carlo simulation study on risk measures estimation in time series derived from FIEGARCH process. Prass and Lopes (2011b) analyze a portfolio composed by stocks from the Brazilian market Bovespa. The authors consider the econometric approach to estimate the risk measure VaR and use FIEGARCH models to obtain the conditional variance of the time series. We also present here the h step-ahead forecast, for $h \in \{1, \cdots, 50\}$, for the conditional variance of simulated FIEGARCH processes.

4.1 Data Generating Process

In the following, the notation Mi, for $i \in \{1, \cdots, 5\}$, indicates the simulated FIEGARCH(p, d, q) models. For each model we consider $re = 1,000$ replications, with sample size $n \in \{2,000; 5,000\}$. The value $n = 2,000$ is the approximated size of the observed time series considered in Prass and Lopes (2011b). The value $n = 5,000$ was chosen to analyze the estimators asymptotic properties.

For each $i \in \{1, \cdots, 5\}$, $\{X_{i,t}\}_{t=1}^{n}$ denotes any time series, with sample size n, derived from the model Mi. To generate these time series we set $Z_{i,0} \sim \mathcal{N}(0, 1)$, for all $i \in \{1, \cdots, 5\}$, consider the expressions given in Definition 2.1 and the equality given in expression (2.8). Then, for each $i \in \{1, \cdots, 5\}$,

$$\begin{aligned} X_{i,t} &= \sigma_{i,t} Z_{i,t} \\ \ln(\sigma_{i,t}^2) &= \omega + \frac{\alpha(\mathcal{B})}{\beta(\mathcal{B})}(1-\mathcal{B})^{-d} g(Z_{i,t-1}) = \sum_{k=0}^{\infty} \lambda_{d,k} g(Z_{i,t-1}) \approx \sum_{k=0}^{m} \lambda_{d,k} g(Z_{i,t-1}), \end{aligned}$$

for m large enough. We set $m = 50,000$ based on the result presented in Theorem 2.1. The parameters value for each model are given in Table 4.1.

Table 4.1: Parameters value for the models.

Model	Parameter									
	d	θ	γ	ω	α_1	α_2	β_1	β_2	β_3	β_4
M1: FIEGARCH$(2, d, 1)$	0.4495	-0.1245	0.3662	-6.5769	-1.1190	-0.7619	-0.6195	-	-	-
M2: FIEGARCH$(0, d, 4)$	0.2391	-0.0456	0.3963	-6.6278	-	-	0.2289	0.1941	0.4737	-0.4441
M3: FIEGARCH$(0, d, 1)$	0.4312	-0.1095	0.3376	-6.6829	-	-	0.5454	-	-	-
M4: FIEGARCH$(0, d, 1)$	0.3578	-0.1661	0.2792	-7.2247	-	-	0.6860	-	-	-
M5: FIEGARCH$(1, d, 1)$	0.4900	-0.0215	0.3700	-5.8927	0.1409	-	-0.1611	-	-	-

From Theorem 2.1, we have

$$\lambda_{d,k} \sim \frac{1}{\Gamma(d) k^{1-d}} \frac{\alpha(1)}{\beta(1)}, \quad \text{as } k \to \infty,$$

and we conclude that $\lambda_{d,k} = o(k^d)$ and $\lambda_{d,k} = O(k^{d-1})$, as k goes to infinity. However, the speed of the convergence varies from model to model, as we show in Table 4.2. For simplicity, let $Q_1(\cdot)$ and $Q_2(\cdot)$ be defined as

$$Q_1(k) := \frac{\lambda_{d,k}}{k^d} \quad \text{and} \quad Q_2(k) := \lambda_{d,k}\Big(\frac{1}{\Gamma(d) k^{1-d}} \frac{\alpha(1)}{\beta(1)}\Big)^{-1}, \quad \text{for all } k > 0.$$

Table 4.2 presents the values of the coefficients $\lambda_{d,k}$, given in Proposition 2.1, for $k \in \{0; 10;$ $100;\ 1{,}000;\ 5{,}000;\ 10{,}000;\ 20{,}000;\ 50{,}000;\ 100{,}000\}$, for each simulated model Mi, $i \in \{1, \cdots, 5\}$. Note that, for $k \geq 5{,}000$, the coefficient values decrease slowly. We also report in Table 4.2 $Q_1(k)$ and $Q_2(k)$ values for the correspondent $\lambda_{d,k}$ value. Note that, for $k \in \{10{,}000;\ 50{,}000;\ 100{,}000\}$, the value $Q_1(k)$ is very close to zero, for all models. Also notice that, while $Q_2(k)$ converges to 1 faster for model M1 than for the other models.

Table 4.2: Coefficients $\lambda_{d,k}$ and the quotients $Q_1(k)$ and $Q_2(k)$, for different values of k, for all models.

Model	M1			M2			M3			M4			M5		
k	$\lambda_{d,k}$	$Q_1(k)$	$Q_2(k)$	$\lambda_{d,k}$	$Q_1(k)$	$Q_2(k)$	$\lambda_{d,k}$	$Q_1(k)$	$Q_2(k)$	$\lambda_{d,k}$	$Q_1(k)$	$Q_2(k)$	$\lambda_{d,k}$	$Q_1(k)$	$Q_2(k)$
0	1.00000	-	-	1.00000	-	-	1.00000	-	-	1.00000	-	-	1.00000	-	-
10	0.26537	0.09426	1.04410	-0.09039	-0.05212	-1.08434	0.31434	0.11647	1.08789	0.36874	0.16178	1.26414	0.12291	0.03977	0.97189
100	0.07167	0.00904	1.00173	0.01450	0.00482	1.00292	0.07844	0.01077	1.00576	0.06738	0.01297	1.01350	0.03897	0.00408	0.99720
1,000	0.02015	0.00090	1.00017	0.00251	0.00048	1.00027	0.02106	0.00107	1.00056	0.01517	0.00128	1.00129	0.01207	0.00041	0.99972
5,000	0.00830	0.00018	1.00003	0.00074	0.00010	1.00005	0.00843	0.00021	1.00011	0.00539	0.00026	1.00026	0.00531	0.00008	0.99994
10,000	0.00567	0.00009	1.00002	0.00043	0.00005	1.00003	0.00568	0.00011	1.00006	0.00345	0.00013	1.00013	0.00373	0.00004	0.99997
25,000	0.00342	0.00004	1.00001	0.00022	0.00002	1.00001	0.00337	0.00004	1.00002	0.00192	0.00005	1.00005	0.00234	0.00002	0.99999
50,000	0.00234	0.00002	1.00000	0.00013	0.00001	1.00001	0.00227	0.00002	1.00001	0.00123	0.00003	1.00003	0.00164	0.00001	0.99999
100,000	0.00160	0.00001	1.00000	0.00008	0.00000	1.00000	0.00153	0.00001	1.00001	0.00079	0.00001	1.00001	0.00115	0.00000	1.00000

4.2 Estimation

In this study we consider the quasi-likelihood method to estimate the parameters of FIEGARCH models for the simulated time series. Given any time series $\{X_t\}_{t=1}^n$, this method assumes that $X_t|\mathcal{F}_{t-1}$, for all $t \in \mathbb{Z}$, is normally distributed. The vector of unknown parameters is denoted by

$$\boldsymbol{\eta} = (d; \omega; \theta; \lambda; \alpha_1, \cdots, \alpha_p; \beta_1, \cdots, \beta_q)' \ \in \mathbb{R}^{p+q+4}$$

and the estimator $\hat{\boldsymbol{\eta}}$ of $\boldsymbol{\eta}$ is the value that maximizes

$$\ln\big(\ell(\boldsymbol{\theta}; \mathbf{x})\big) = -\frac{n}{2}\ln(2\pi) - \frac{1}{2}\sum_{t=1}^{n}\left[\ln(\sigma_t^2) + \frac{X_t^2}{\sigma_t^2}\right]. \tag{4.1}$$

Since the processes $\{X_t\}_{t<1}$ and $\{Z_t\}_{t<1}$ are unknown, we need to consider a set I_0 of initial conditions in order to start the recursion and to obtain the random variable $\ln(\sigma_t^2)$, for $t \in \{1, \cdots, n\}$. Then we use these estimated values to solve (4.1). For this simulation study we assume, as initial conditions, $Z_t = 0$, $\sigma_t^2 = \hat{\sigma}_X^2$ and $X_t := \sigma_t Z_t = 0$, whenever $t < 1$, where $\hat{\sigma}_X^2$ is the sample variance of $\{X_t\}_{t=1}^n$. This is the initial set suggested by Bollerslev and Mikkelsen (1996). The random variables $\ln(\sigma_t^2)$, for $t \in \{1, \cdots, n\}$, are then estimated upon considering the set I_0 of initial conditions and the known values $\{X_t\}_{t=1}^n$. The infinite sum in the polynomial $\lambda(\cdot)$ is truncated at $m = n$, where n is the available sample size.

Figures 4.1 - 4.5 present the kernel distribution of the parameter estimators for each considered model when $n \in \{2,000; 5,000\}$. These graphs help to illustrate the results presented in Table 4.3. It is easy to see that, for most estimates, the density function is approximately symmetric. For some parameters, we notice the presence of possible outliers, see for instance the graphs for parameter d, from model M1, and parameters β_i, with $i \in \{2, 3, 4\}$, for model M2. Although the graphs for $n = 2,000$ and $n = 5,000$ are similar, one observes that, as expected, the observations tend to concentrate closer to the mean when $n = 5,000$.



Figure 4.1: Kernel density function of the estimates for model M1, for $n \in \{2,000;\ 5,000\}$.



Figure 4.2: Kernel density function of the estimates for model M2, for $n \in \{2,000;\ 5,000\}$.



Figure 4.3: Kernel density function of the estimates for model M3, for $n \in \{2,000;\ 5,000\}$.



Figure 4.4: Kernel density function of the estimates for model M4, for $n \in \{2,000;\ 5,000\}$.



Figure 4.5: Kernel density function of the estimates for model M5, for $n \in \{2,000;\ 5,000\}$.

For any model, $\hat{\eta}_k$ denotes the η estimate in the k-th replication, where $k \in \{1, \cdots, re\}$, $re = 1,000$ and η is any vector parameter given in Table 4.1. Table 4.3 summarizes the results on the parameter estimation procedure. The reported values are the mean $\bar{\eta}$, the standard deviation (sd), the bias ($bias$), the mean absolute error (mae) and the mean square error (mse) values, defined by

$$\bar{\eta} := \frac{1}{re}\sum_{k=1}^{re}\hat{\eta}_k, \quad sd := \sqrt{\frac{1}{re}\sum_{k=1}^{re}(\hat{\eta}_k - \bar{\eta})^2}, \quad bias := \frac{1}{re}\sum_{k=1}^{re} e_k, \quad mae := \frac{1}{re}\sum_{k=1}^{re}|e_k|, \quad \text{and} \quad mse := \frac{1}{re}\sum_{k=1}^{re} e_k^2,$$

where $e_k := \hat{\eta}_k - \eta$, for $k \in \{1, \cdots, re\}$.

From Table 4.3 we conclude that, given the models complexity, the quasi-likelihood method performs relatively well. Since model M2 presents more parameters than the other models, which implies a higher dimension maximization problem, one would expect that the quasi-likelihood method would present the worst performance in this case. However, the estimation results for model M2 ($d = 0.2391$) and M4 ($d = 0.3578$) are similar and the quasi-likelihood method performs better for model M2 than for models M1 ($d = 0.4495$), M3 ($d = 0.4312$) and M5 ($d = 0.4900$). By comparing the mae and mse values, given in Table 4.3, we conclude that the worst performance occurs for models M1 and M5. This outcome is explained by the fact that the parameter d is very close to the non-stationary region for model M5 and, for model M1, not only $p = 2$ but also $d = 0.4495$, which implies a more complex model with stronger long-range dependence. The small $bias$ values indicate that, for all parameters, the mean estimated value is very close to the true value. Although for $n = 2,000$ the standard deviation of several estimates is high if compared with the mean estimated value, as expected, the estimators performance improves as the sample size increases.

Table 4.3: Estimation results.

Sample Size (n)		$2,000$					$5,000$				
Model	η	$\bar{\eta}$	*sd*	*bias*	*mae*	*mse*	$\bar{\eta}$	*sd*	*bias*	*mae*	*mse*
M1	$d = 0.4495$	0.4104	0.0743	-0.0391	0.0522	0.0070	0.4133	0.0776	-0.0362	0.0495	0.0073
	$\theta = -0.1245$	-0.1255	0.0230	-0.0010	0.0185	0.0005	-0.1255	0.0220	-0.0010	0.0176	0.0005
	$\gamma = 0.3662$	0.3651	0.0480	-0.0011	0.0387	0.0023	0.3653	0.0458	-0.0009	0.0365	0.0021
	$\omega = -6.5769$	-6.5757	0.3955	0.0012	0.3156	0.1565	-6.5765	0.3916	0.0004	0.3128	0.1534
	$\alpha_1 = -1.1190$	-1.0211	0.3776	0.0979	0.2719	0.1521	-1.0273	0.3664	0.0917	0.2588	0.1427
	$\alpha_2 = -0.7619$	-0.7304	0.3097	0.0315	0.2048	0.0969	-0.7279	0.3035	0.0340	0.1965	0.0933
	$\beta_1 = -0.6195$	-0.4659	0.3784	0.1536	0.2404	0.1668	-0.4766	0.3719	0.1429	0.2282	0.1587
M2	$d = 0.2391$	0.2086	0.1188	-0.0305	0.0959	0.0150	0.2208	0.0833	-0.0183	0.0615	0.0073
	$\theta = -0.0456$	-0.0461	0.0239	-0.0005	0.0192	0.0006	-0.0459	0.0148	-0.0003	0.0117	0.0002
	$\gamma = 0.3963$	0.3933	0.0497	-0.0030	0.0383	0.0025	0.3949	0.0323	-0.0014	0.0250	0.0010
	$\omega = -6.6278$	-6.6325	0.0918	-0.0047	0.0731	0.0085	-6.6271	0.0663	0.0007	0.0518	0.0044
	$\beta_1 = 0.2289$	0.2525	0.1479	0.0236	0.0988	0.0224	0.2422	0.0978	0.0133	0.0631	0.0097
	$\beta_2 = 0.1941$	0.1877	0.1082	-0.0064	0.0656	0.0117	0.1938	0.0819	-0.0003	0.0421	0.0067
	$\beta_3 = 0.4737$	0.4605	0.1204	-0.0132	0.0631	0.0147	0.4655	0.0859	-0.0082	0.0398	0.0074
	$\beta_4 = -0.4441$	-0.4535	0.1302	-0.0094	0.0843	0.0170	-0.4483	0.0732	-0.0042	0.0489	0.0054
M3	$d = 0.4312$	0.3703	0.1474	-0.0609	0.1040	0.0254	0.4148	0.0577	-0.0164	0.0458	0.0036
	$\theta = -0.1095$	-0.1121	0.0227	-0.0026	0.0180	0.0005	-0.1100	0.0142	-0.0005	0.0113	0.0002
	$\gamma = 0.3376$	0.3372	0.0433	-0.0004	0.0347	0.0019	0.3366	0.0272	-0.0010	0.0218	0.0007
	$\omega = -6.6829$	-6.6954	0.4058	-0.0125	0.3253	0.1648	-6.6942	0.3516	-0.0113	0.2865	0.1238
	$\beta_1 = 0.5454$	0.5839	0.1508	0.0385	0.1203	0.0242	0.5582	0.0782	0.0128	0.0625	0.0063
M4	$d = 0.3578$	0.3094	0.1240	-0.0484	0.0921	0.0177	0.3408	0.0590	-0.0170	0.0453	0.0038
	$\theta = -0.1661$	-0.1698	0.0207	-0.0037	0.0167	0.0004	-0.1667	0.0136	-0.0006	0.0108	0.0002
	$\gamma = 0.2792$	0.2794	0.0368	0.0002	0.0293	0.0014	0.2787	0.0227	-0.0005	0.0182	0.0005
	$\omega = -7.2247$	-7.2171	0.2729	0.0076	0.2150	0.0745	-7.2176	0.2204	0.0071	0.1750	0.0486
	$\beta_1 = 0.6860$	0.7095	0.1003	0.0235	0.0802	0.0106	0.6963	0.0570	0.0103	0.0461	0.0034
M5	$d = 0.4900$	0.4390	0.1602	-0.0510	0.1216	0.0283	0.4589	0.1092	-0.0311	0.0765	0.0129
	$\theta = -0.0215$	-0.0228	0.0314	-0.0013	0.0247	0.0010	-0.0214	0.0188	0.0001	0.0150	0.0004
	$\gamma = 0.3700$	0.3704	0.0500	0.0004	0.0401	0.0025	0.3702	0.0334	0.0002	0.0268	0.0011
	$\omega = -5.8927$	-5.9000	0.2418	-0.0073	0.1926	0.0585	-5.8988	0.2137	-0.0061	0.1705	0.0457
	$\alpha_1 = 0.1409$	0.1605	0.4387	0.0196	0.3619	0.1929	0.1944	0.3916	0.0535	0.3157	0.1562
	$\beta_1 = -0.1611$	-0.0817	0.4680	0.0794	0.3762	0.2253	-0.0608	0.4370	0.1003	0.3311	0.2010

4.3 Forecasting

In what follows we discuss the simulation results related to forecasting based on the fitted FIEGARCH models. To access the models forecast performance, during the generating process, we create 50 extra values for each simulated time series. Those values are used here to compare with the h-step ahead forecast, for $h \in \{1, \cdots, 50\}$. For each replication, the h-step ahead predictor for σ^2_{n+h}, for all $h \in \{1, \cdots, 50\}$, is obtained from expression (3.7) upon replacing the true parameters values $\boldsymbol{\eta} = (d; \omega; \theta; \lambda; \alpha_1, \cdots, \alpha_p; \beta_1, \cdots, \beta_q)'$ by the estimated ones. Based on the fact that $\mathbb{E}(X^2_{n+h}|\mathcal{F}_n) = \mathbb{E}(\sigma^2_{n+h}|\mathcal{F}_n)$, we also set $\tilde{X}^2_{n+h} := \tilde{\sigma}^2_{n+h}$, for all $h > 0$.

Table 4.4 presents the mean over $1,000$ simulated values of $\sigma^2_{i,n+h}$ and $X^2_{i,n+h}$, obtained from model Mi, for each $i \in \{1, \cdots, 5\}$, the corresponding h-step ahead predicted values $\tilde{\sigma}^2_{i,t+h} := \tilde{X}^2_{i,t+h}$ and, in parenthesis, the mean absolute error of forecast (*mae*), for $h \in \{1, \cdots, 10\}$ and forecasting origin $n \in \{2,000; 5,000\}$. Figure 4.6 illustrates the results given in Table 4.4.

From Table 4.4 we conclude that, for $\sigma^2_{i,n+h}$, $h \in \{1, \cdots, 10\}$ and $i \in \{1, \cdots, 5\}$, the predicted values are relatively close to the simulated ones, which is indicated by the small *mae* values. We also observe that the *mae* value increases as h increases. This result is expected and it is theoretically explained in Proposition 3.1. Table 4.4 also shows that, for $X^2_{i,n+h}$, the *mae* is usually high, if compared to the mean simulated and mean predicted values. Therefore, we conclude that $\tilde{X}^2_{n+h} :=$

Table 4.4: Mean simulated values for $\sigma^2_{i,n+h}$ and $X^2_{i,n+h}$, obtained from model Mi, $i \in \{1, \cdots, 5\}$, and the mean predicted values $\tilde{\sigma}^2_{i,n+h} = \tilde{X}^2_{i,n+h}$, for $h \in \{1, \cdots, 10\}$ and forecasting origin $n \in \{2,000; 5,000\}$. In parenthesis, the mean absolute error of forecast.

n	2,000			5,000		
h	Predictor	$\sigma^2_{i,n+h}$	$X^2_{i,n+h}$	Predictor	$\sigma^2_{i,n+h}$	$X^2_{i,n+h}$
M1 ($i = 1$)						
1	0.00183	0.00182 (0.00014)	0.00165 (0.00169)	0.00181	0.00180 (0.00013)	0.00163 (0.00165)
2	0.00185	0.00182 (0.00040)	0.00197 (0.00179)	0.00178	0.00172 (0.00036)	0.00167 (0.00172)
3	0.00180	0.00179 (0.00054)	0.00186 (0.00185)	0.00179	0.00176 (0.00053)	0.00185 (0.00188)
4	0.00181	0.00185 (0.00068)	0.00194 (0.00195)	0.00176	0.00177 (0.00067)	0.00174 (0.00175)
5	0.00179	0.00185 (0.00072)	0.00190 (0.00195)	0.00177	0.00179 (0.00068)	0.00162 (0.00170)
6	0.00180	0.00184 (0.00075)	0.00181 (0.00178)	0.00176	0.00176 (0.00073)	0.00148 (0.00161)
7	0.00179	0.00182 (0.00073)	0.00175 (0.00183)	0.00177	0.00172 (0.00068)	0.00166 (0.00172)
8	0.00179	0.00181 (0.00075)	0.00179 (0.00185)	0.00176	0.00173 (0.00071)	0.00177 (0.00181)
9	0.00179	0.00178 (0.00074)	0.00183 (0.00186)	0.00177	0.00176 (0.00070)	0.00173 (0.00180)
10	0.00179	0.00181 (0.00076)	0.00183 (0.00195)	0.00176	0.00175 (0.00069)	0.00188 (0.00191)
M2 ($i = 2$)						
1	0.00145	0.00145 (0.00011)	0.00148 (0.00148)	0.00144	0.00144 (0.00007)	0.00139 (0.00143)
2	0.00147	0.00147 (0.00032)	0.00143 (0.00144)	0.00143	0.00143 (0.00031)	0.00147 (0.00145)
3	0.00143	0.00143 (0.00034)	0.00145 (0.00137)	0.00144	0.00144 (0.00033)	0.00158 (0.00151)
4	0.00144	0.00146 (0.00036)	0.00144 (0.00139)	0.00143	0.00145 (0.00037)	0.00153 (0.00145)
5	0.00144	0.00144 (0.00042)	0.00135 (0.00138)	0.00143	0.00144 (0.00043)	0.00148 (0.00145)
6	0.00141	0.00141 (0.00041)	0.00142 (0.00147)	0.00143	0.00146 (0.00043)	0.00140 (0.00136)
7	0.00143	0.00144 (0.00043)	0.00150 (0.00147)	0.00142	0.00143 (0.00041)	0.00142 (0.00141)
8	0.00142	0.00143 (0.00043)	0.00146 (0.00144)	0.00143	0.00143 (0.00042)	0.00142 (0.00144)
9	0.00142	0.00143 (0.00044)	0.00131 (0.00135)	0.00142	0.00143 (0.00044)	0.00143 (0.00141)
10	0.00143	0.00142 (0.00042)	0.00143 (0.00144)	0.00142	0.00142 (0.00042)	0.00138 (0.00142)
M3 ($i = 3$)						
1	0.00158	0.00157 (0.00009)	0.00167 (0.00158)	0.00151	0.00150 (0.00005)	0.00147 (0.00139)
2	0.00157	0.00158 (0.00033)	0.00152 (0.00147)	0.00151	0.00150 (0.00027)	0.00149 (0.00147)
3	0.00157	0.00159 (0.00042)	0.00145 (0.00154)	0.00151	0.00151 (0.00041)	0.00151 (0.00150)
4	0.00157	0.00157 (0.00049)	0.00155 (0.00156)	0.00151	0.00151 (0.00048)	0.00151 (0.00152)
5	0.00157	0.00157 (0.00055)	0.00161 (0.00166)	0.00152	0.00151 (0.00050)	0.00153 (0.00150)
6	0.00157	0.00158 (0.00060)	0.00152 (0.00154)	0.00152	0.00151 (0.00053)	0.00147 (0.00153)
7	0.00158	0.00157 (0.00060)	0.00151 (0.00152)	0.00152	0.00152 (0.00057)	0.00149 (0.00156)
8	0.00158	0.00157 (0.00060)	0.00170 (0.00178)	0.00152	0.00152 (0.00061)	0.00150 (0.00153)
9	0.00158	0.00160 (0.00064)	0.00170 (0.00170)	0.00152	0.00150 (0.00059)	0.00148 (0.00147)
10	0.00159	0.00162 (0.00066)	0.00173 (0.00176)	0.00152	0.00151 (0.00059)	0.00143 (0.00147)
M4 ($i = 4$)						
1	0.00098	0.00097 (0.00006)	0.00091 (0.00089)	0.00092	0.00092 (0.00003)	0.00089 (0.00085)
2	0.00097	0.00096 (0.00019)	0.00098 (0.00094)	0.00092	0.00092 (0.00018)	0.00088 (0.00085)
3	0.00096	0.00095 (0.00028)	0.00098 (0.00097)	0.00092	0.00092 (0.00026)	0.00091 (0.00089)
4	0.00095	0.00096 (0.00034)	0.00104 (0.00100)	0.00092	0.00092 (0.00031)	0.00090 (0.00091)
5	0.00094	0.00098 (0.00039)	0.00102 (0.00102)	0.00092	0.00092 (0.00035)	0.00095 (0.00093)
6	0.00094	0.00099 (0.00044)	0.00096 (0.00098)	0.00092	0.00093 (0.00037)	0.00080 (0.00083)
7	0.00094	0.00098 (0.00044)	0.00101 (0.00103)	0.00092	0.00092 (0.00038)	0.00101 (0.00095)
8	0.00093	0.00098 (0.00045)	0.00102 (0.00106)	0.00091	0.00093 (0.00041)	0.00093 (0.00099)
9	0.00093	0.00098 (0.00047)	0.00103 (0.00102)	0.00091	0.00092 (0.00041)	0.00092 (0.00097)
10	0.00093	0.00098 (0.00047)	0.00103 (0.00103)	0.00091	0.00094 (0.00044)	0.00099 (0.00101)
M5 ($i = 5$)						
1	0.00292	0.00292 (0.00018)	0.00310 (0.00296)	0.00298	0.00296 (0.00012)	0.00277 (0.00276)
2	0.00295	0.00297 (0.00059)	0.00301 (0.00290)	0.00295	0.00291 (0.00056)	0.00298 (0.00299)
3	0.00295	0.00296 (0.00059)	0.00293 (0.00291)	0.00296	0.00294 (0.00058)	0.00315 (0.00292)
4	0.00295	0.00295 (0.00061)	0.00294 (0.00288)	0.00295	0.00298 (0.00061)	0.00322 (0.00308)
5	0.00295	0.00296 (0.00061)	0.00296 (0.00284)	0.00295	0.00298 (0.00063)	0.00310 (0.00292)
6	0.00295	0.00296 (0.00061)	0.00298 (0.00295)	0.00296	0.00299 (0.00061)	0.00321 (0.00304)
7	0.00295	0.00296 (0.00064)	0.00303 (0.00289)	0.00295	0.00300 (0.00063)	0.00277 (0.00285)
8	0.00295	0.00296 (0.00061)	0.00306 (0.00298)	0.00295	0.00293 (0.00061)	0.00307 (0.00307)
9	0.00295	0.00297 (0.00065)	0.00289 (0.00297)	0.00295	0.00298 (0.00066)	0.00304 (0.00302)
10	0.00295	0.00294 (0.00063)	0.00297 (0.00309)	0.00295	0.00296 (0.00064)	0.00308 (0.00299)

$\tilde{\sigma}^2_{n+h}$ is a poor estimator for X^2_{n+h}. This result is not a surprise since the main purpose of FIEGARCH models is to estimate the logarithm of the conditinal variance of the process and not the process itself.

As expected, in all cases, the models' forecasting performance improves as n increases. Notice, however, that the difference in the *mae* values, from $n = 2,000$ to $n = 5,000$, is small. This is so because the coefficients $\lambda_{d,k}$ converges to zero, as k goes to infinity. Therefore, it is expected that, for some $m \in \mathbb{N}$ and any $M > 0$, using the last m or the last $m + M$ known values to calculate the h-step ahead forecast value for the process will not considerably change the results.



Figure 4.6: Mean simulated values for $\sigma^2_{i,n+h}$ and $X^2_{i,n+h}$, obtained from model Mi, $i \in \{1, \cdots, 5\}$, and mean predicted values $\tilde{\sigma}^2_{i,n+h} = \tilde{X}^2_{i,n+h}$, for $h \in \{1, \cdots, 10\}$ and forecasting origin $n \in \{2,000; 5,000\}$. For each model the graphs correspond to the mean taken over 1,000 replications.

Figure 4.7 shows the mean simulated values of $\sigma^2_{i,n+h}$ and $X^2_{i,n+h}$, for $i \in \{1, \cdots, 5\}$, and the corresponding predicted values, obtained from expression (3.7), for $h \in \{1, \cdots, 50\}$ and forecasting origin $n \in \{2,000; 5,000\}$.



Figure 4.7: Graphs of the h-step ahead forecast values $\tilde{X}^2_{i,n+h} = \tilde{\sigma}^2_{i,n+h}$, for $h \in \{1, \cdots, 50\}$ and forecasting origin $n \in \{2,000; 5,000\}$, and the corresponding simulated values $X^2_{i,n+h}$ and $\sigma^2_{i,n+h}$ from model Mi, $i \in \{1, \cdots, 5\}$. For each model the graphs correspond to the mean taken over 1,000 replications.

From Figure 4.7 we observe that, for all models, the predicted value converges to a constant as h increases. This is expected since the h-step ahead predictor is defined in terms of the conditional expectation. We also observe that, if we had considered the h-step ahead predictor $\breve{\sigma}^2_{i,n+h}$ instead of $\tilde{\sigma}^2_{i,n+h}$, the estimation bias would have been much higher. Indeed, from expression (3.5), $\breve{\sigma}^2_{i,n+h}$ converges to $e^{\omega(i)}$ as h goes to infinity, where $\omega(i)$ denotes the parameter ω for model Mi, for each $i \in \{1, \cdots, 5\}$. The values of $\omega(i)$ are given in Table 4.1 and the values of $e^{\omega(i)}$, for $i \in \{1, \cdots, 5\}$, are respectively 0.0014, 0.0013, 0.0013, 0.0007 and 0.0028.

Notice now that equations (3.5) and (3.8) imply

$$\tilde{\sigma}^2_{i,n+h} := \check{\sigma}^2_{i,n+h}\Big(1+\frac{1}{2}\sigma_g^2\sum_{k=0}^{h-2}\lambda^2_{d,k}\Big) \overset{h\to\infty}{\longrightarrow} e^{\omega(i)}\Big(1+\frac{1}{2}\sigma_g^2(i)\sum_{k=0}^{\infty}\lambda^2_{d,k}(i)\Big) \approx e^{\omega(i)}\Big(1+\frac{1}{2}\sigma_g^2(i)\sum_{k=0}^{m}\lambda^2_{d,k}(i)\Big), \quad (4.2)$$

for any $i \in \{1, \cdots, 5\}$ and m sufficiently large. By taking $m = 50{,}000$ in (4.2), the limits are 0.00173, 0.00142, 0.00155, 0.00091 and 0.00294, respectively, for $i \in \{1, \cdots, 5\}$. Upon comparing these values with the limiting constants in Figure 4.7, we conclude that, in fact, expression (4.2) holds for all models.

5 Analysis of an Observed Time Series

This section presents the analysis of the São Paulo Stock Exchange Index (Bovespa Index or IBovespa) log-return time series. We consider the FIEGARCH model, fully described in this paper, and we compare its forecasting performance with other ARCH-type models. The total number of observations for the IBovespa time series is $n = 1737$. We consider the first 1717 observations to fit the models and we reserve the last 20 ones to compare with the out-of-sample forecast.

Figure 5.1 (a) presents IBovespa time series $\{P_t\}_{t=1}^{1718}$, in the period of January/1995 to December/2001. We observe a strong decay in the index value close to $t = 1,000$ (that is, January 15, 1999). This period is characterized by the Real (the Brazilian currency) devaluation. Figures 5.1 (b) and (c) present, respectively, the IBovespa log-return time series, $\{r_t\}_{t=1}^{1717}$, and the square of the log-return time series, $\{r_t^2\}_{t=1}^{1717}$, in the same period. Observe that the log-return series presents the stylized facts of financial time series such as apparent stationarity, mean around zero and clusters of volatility. Also, in Figure 5.2 we observe that, while the log-return series presents almost no correlation, the sample autocorrelation of the square of the log-return series assumes high values for several lags, pointing to the existence of heteroskedasticity and possibly long memory. Notice that the periodogram of $\{\ln(r_t^2)\}_{t=1}^{1717}$, presented in Figure 2.3 (c), also indicates possibly long-memory in the conditional variance. Regarding the histogram and the QQ-Plot, we observe that the distribution of the log-return series seems approximately symmetric and leptokurtic.



(a) $\{P_t\}_{t=1}^{1718}$ (b) $\{r_t\}_{t=1}^{1717}$ (c) $\{r_t^2\}_{t=1}^{1717}$

Figure 5.1: Time series: (a) Bovespa index; (b) IBovespa log-returns; (c) square of the IBovespa log-returns, in the period of January/1995 to December/2001.

The analysis of the sample autocorrelation function suggests an ARMA(p_1, q_1)-FIEGARCH(p_2, d, q_2) model. While an ARMA model accounts for the correlation among the log-returns, a FIEGARCH model take into account the long memory (in the conditional variance) and the heteroskedasticity characteristics of the time series.

Initially, we consider all possible models with $p_1, q_1 \in \{0, 1, 2, 3\}$ and $p_2, q_2 \in \{0, 1, 2\}$ and applied the quasi-likelihood method to estimate the unknown parameters. Then, we select the best models,



(a) (b) (c) (d)

Figure 5.2: (a) Histogram; (b) QQ-Plot and (c) sample autocorrelation of the IBovespa log-return series and (d) sample autocorrelation of the square of the IBovespa log-return series.

with respect to the log-likelihood, AIC, BIC and HQC criteria. After removing the non-significant parameters and performing the residual analysis we conclude that the best model is an ARMA(0,1)-FIEGARCH$(0, d, 1)$. We compare the forecasting performance of this model with other ARCH-type models.

Table 5.1: Fitted Models and their respective log-likelihood values.

Estimate	ARMA(0,1) + ARCH(1)	ARMA(0,1) + ARCH(6)	ARMA(0,1) + GARCH(1,1)	ARMA(0,1) + EGARCH(0,1)	ARMA(0,1) + FIEGARCH(0,d,1)
$\hat{\theta}_1$	-0.1138 (0.0200)	-0.0642 (0.0267)	-0.0647 (0.0266)	-0.0751 (0.0254)	-0.0776 (0.0257)
$\hat{\omega}$	0.0004 (0.0000)	0.0002 (0.0000)	0.0000 (0.0000)	-7.4694 (0.0969)	-7.2247 (0.2143)
$\hat{\alpha}_1$	0.6071 (0.0581)	0.2307 (0.0417)	0.2019 (0.0247)	-	-
$\hat{\alpha}_2$	-	0.1540 (0.0333)	-	-	-
$\hat{\alpha}_3$	-	0.1852 (0.0390)	-	-	-
$\hat{\alpha}_4$	-	0.1145 (0.0348)	-	-	-
$\hat{\alpha}_5$	-	0.0641 (0.0290)	-	-	-
$\hat{\alpha}_6$	-	0.0635 (0.0257)	-	-	-
$\hat{\beta}_1$	-	-	0.7659 (0.0271)	0.9373 (0.0103)	0.6860 (0.0986)
$\hat{d}$	-	-	-	-	0.3578 (0.0810)
$\hat{\theta}$	-	-	-	-0.1653 (0.0197)	-0.1661 (0.0224)
$\hat{\gamma}$	-	-	-	0.2782 (0.0300)	0.2972 (0.0332)
log-likelihood	3934.337	4060.372	4072.622	4137.6251	4138.5518

Note: The number in parenthesis corresponds to the standard error of the estimate.

The estimated coefficients for each model are given in Table 5.1, with the corresponding log-likelihood value. Notice that, the FIEGARCH model fitted to this time series present the same parameters values as model M4 considered in the simulated study in Section 4. Table 5.2 presents the mean absolute error (mae), the mean percentage error (mpe) and the maximum absolute error (max_{ae}) of forecast, respectively defined as

$$mae = \frac{1}{20}\sum_{h=1}^{20}|e_{n+h}|, \quad mpe := \frac{1}{20}\sum_{h=1}^{20}\frac{|e_{n+h}|}{r_{n+h}^2} \quad \text{and} \quad max_{ae} := \max_{h\in\{1,\cdots,20\}}\{|e_{n+h}|\}$$

where, $e_{n+h} := \hat{r}_{n+h}^2 - r_{n+h}^2$, for $h \in \{1, \cdots, 20\}$ and $n = 1717$, is the h-step ahead forecast error. Note that, from the ARMA(0,1) part of the models, $r_t = X_t - \theta_1 X_{t-1}$, where $X_t = \sigma_t Z_t$, for all $t \in \mathbb{Z}$. Since we define $\hat{r}_{t+h}^2 = \mathbb{E}(r_{t+h}^2|\mathcal{F}_t)$ and σ_t^2 is $\mathcal{F}_{t-1}$-measurable, for all $t \in \mathbb{Z}$, by elementary calculations we conclude that, $\hat{r}^2_{n+1} = \sigma^2_{n+1} + \theta_1^2 X_n^2$ and $\hat{r}^2_{n+h} = \hat{\sigma}^2_{n+h} + \theta_1^2 \hat{\sigma}^2_{n+h-1}$, for all $h > 1$, with $\hat{\sigma}^2_{n+1} = \sigma^2_{n+1}$. For EGARCH and FIEGARCH models, $\hat{\sigma}^2_{n+1}$ is replaced by $\tilde{\sigma}^2_{n+1}$, given in expression (3.7), and $\breve{\sigma}^2_{n+h} := \exp\{\hat{\ln}(\sigma^2_{n+h})\}$, where $\hat{\ln}(\sigma^2_{n+h})$ is defined in Proposition 3.1.

Table 5.2: Mean absolute error (*mae*), mean percentage error (*mpe*) and maximum absolute error (max_{ae}) of forecasting for the models in Table 5.1.

Model	ARMA(0,1) + ARCH(1)	ARMA(0,1) + ARCH(6)	ARMA(0,1) + GARCH(1,1)	ARMA(0,1) + EGARCH(1,1)		ARMA(0,1) + FIEGARCH(1,d,1)	
Predictor	$\hat{\sigma}^2_{t+h}$	$\hat{\sigma}^2_{t+h}$	$\hat{\sigma}^2_{t+h}$	$\tilde{\sigma}^2_{t+h}$	$\breve{\sigma}^2_{t+h}$	$\tilde{\sigma}^2_{t+h}$	$\breve{\sigma}^2_{t+h}$
mae	0.00053	0.00045	0.00043	0.00045	0.00044	0.00045	0.00043
mpe	109.40844	68.97817	60.29677	71.33057	61.26625	68.42884	59.88066
max_{ae}	0.00094	0.00094	0.00094	0.00082	0.00087	0.00084	0.00088

Note: The high *mpe* values are due to 5 observations close to zero.

From Table 5.2 we conclude that, given its high *mpe* value, the ARMA(0,1)-ARCH(1) does not fit the data well. In fact, the square of the residuals from this model are still correlated and we use the model only for comparison. The ARMA(0,1)-ARCH(6) model performed similar to the ARMA(0,1)-GARCH(1,1) model, in terms of both, *mae* and max_{ae} values, presenting a higher *mpe* value. However, the latter is more parsimonious. Although the log-likelihood value is higher (and the max_{ae} value is smaller) for the ARMA(0,1)-EGARCH(0,1) model, the *mae* and the *mpe* values are smaller for the ARMA(0,1)-GARCH(0,d,1) model. Overall, the ARMA(0,1)-FIEGARCH(0,d,1) performs slightly better than the other models.

The fact that all models present a similar perfomance confirms the following, already known in the literature.

- In practice, ARCH(p) models perform relatively well for most applications.
- GARCH(p, q) models are more parsimonious than the ARCH ones. For instance, notice that similar results were obtained here by considering an ARCH(6) model and a GARCH(1, 1) model.
- For EGARCH(p, q) models the conditional variance is defined in terms of the logarithm function and less (usually none) restrictions have to be imposed during parameter estimation. Moreover, EGARCH models are not necessarily more parsimonious than ARCH/GARCH ones since it also carries information on the returns' asymmetry (θ and γ parameters).
- FIEGARCH(p, d, q) models can describe not only the same characteristics as ARCH, GARCH and EGARCH models do, but also the long-memory in the volatility. Also, the performance of all models will be very similar if the volatility presents high persistence. For instance, notice that for the ARCH(6) model $\alpha_1 + \cdots + \alpha_6 = 0.812$, for the GARCH(1, 1) model $\alpha_1 + \beta_1 = 0.9678$ and for the EGARCH model $\beta_1 = 0.9373$, which imply high persistence in the volatility. Moreover, for the FIEGARCH model, we found $d = 0.3578$ with standard error equal to 0.0810, which indicates that the parameter d is statistically different from zero and thus, there is evidence of long-memory in the volatility.
- Given their definition, it is expected that EGARCH and FIEGARCH models will provide better forecasts for $\ln(\sigma^2_{t+h})$ than for σ^2_{t+h} and, consequently, for X^2_{t+h}.

6 Conclusions

Here we show complete mathematical proofs for the stationarity, the ergodicity, the conditions for the causality and invertibility properties, the autocorrelation and spectral density functions decay and the convergence order for the polynomial coefficients that describe the volatility for any FIEGARCH(p, q, d) process. We prove that if $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH(p, d, q) process and $\mathbb{E}\big(\big[\ln(Z_0^2)\big]^2\big) < \infty$, then $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ is an ARFIMA$(q, d, 0)$ process with correlated innovations. Expressions for the kurtosis and the asymmetry measures of any stationary FIEGARCH(p, d, q) process were also provided.

We also prove that if $\{X_t\}_{t\in\mathbb{Z}}$ is a FIEGARCH(p, d, q) process then, it is a martingale difference with respect to the filtration $\{\mathcal{F}_t\}_{t\in\mathbb{Z}}$, where $\mathcal{F}_t := \sigma(\{Z_s\}_{s\leq t})$. The h-step ahead forecast for the processes $\{X_t\}_{t\in\mathbb{Z}}$, $\{\ln(\sigma_t^2)\}_{t\in\mathbb{Z}}$ and $\{\ln(X_t^2)\}_{t\in\mathbb{Z}}$ are given with their respective mean square error forecast. Since $\mathbb{E}(\sigma_{t+h}^2|\mathcal{F}_t)$ cannot be easily calculated for FIEGARCH models, we also discuss some alternative estimators for the h-step ahead forecast of σ_{t+h}^2, for all $h > 0$.

We present a Monte Carlo simulation study showing how to perform the generation, the estimation and the forecasting of five different FIEGARCH models. The parameter selection of these five models are related to the real time series analyzed in Prass and Lopes (2011a). Parameter estimation was performed by considering the well known quasi-likelihood method. We conclude that, given the complexity of FIEGARCH models, the quasi-likelihood method performs relatively well, which is indicated by the small *bias*, *mae* and *mse* values for the estimates. Regarding the h-step ahead forecast for the processes $\{\sigma_t^2\}_{t\in\mathbb{Z}}$ and $\{X_t^2\}_{t\in\mathbb{Z}}$, we observe that the mean absolute error of forecast decreases as the sample size increases. However, while the conditional variance is well estimated, which is indicated by the small *mae* values, the estimator $\tilde{X}_{n+h}^2 := \tilde{\sigma}_{t+h}^2$, which is an approximation for $\hat{X}_{t+h}^2 := \mathbb{E}(X_{n+h}^2|\mathcal{F}_n) = \hat{\sigma}_{n+h}^2$, does not perform well in predicting X_{n+h}^2. This result is expected since the purpose of the model is to forecast the logarithm of the conditional variance and not the process $\{X_t\}_{t\in\mathbb{Z}}$ itself.

Finally, we present the analysis of the São Paulo Stock Exchange Index (Bovespa Index or IBovespa) log-return time series. We compared the forecasting performance of FIEGARCH models, fully described in this paper, with other ARCH-type models. All models presented a similar performance which was attributed to the fact that the ARCH, GARCH and EGARCH models indicated high persistence in the volatility.

Acknowledgments

S.R.C. Lopes was partially supported by CNPq-Brazil, by CAPES-Brazil, by INCT em *Matemática* and by Pronex *Probabilidade e Processos Estocásticos* - E-26/170.008/2008 -APQ1. T.S. Prass was supported by CNPq-Brazil. The authors are grateful to the (Brazilian) National Center of Super Computing (CESUP-UFRGS) for the computational resources.

References

[1] Baillie, R., T. Bollerslev and H. Mikkelsen (1996). "Fractionally Integrated Generalised Autoregressive Conditional Heteroscedasticity". *Journal of Econometrics*, Vol. **74**, 3-30.

[2] Berkes, I., L. Horváth and P. Kokoszka (2003). "GARCH Processes: Structure and Estimation". *Bernoulli*, Vol. **9**, 201-228.

[3] Berkes, I. and L. Horváth (2003). "The Rate of Consistency of the Quasi-Maximum Likelihood Estimator". *Statistics and Probability Letters*, Vol. **61**, 133-143.

[4] Bloomfield, P. (1985). “On Series Representations for Linear Predictors.” *The Annals of Probability*, Vol. **13**(1), 226-233.

[5] Bollerslev, T. and H.O. Mikkelsen (1996). “Modeling and Pricing Long Memory in Stock Market Volatility”. *Journal of Econometrics*, Vol. **73**, 151-184.

[6] Bondon, P. and W. Palma (2007). “A class of antipersistent processes”. *Journal of Time Series Analysis*, Vol. **28**, 261-273.

[7] Brockwell, P.J. and R.A. Davis (1991). *Time Series: Theory and Methods.* Second Edition. New York: Springer-Verlag.

[8] Doukhan, P., G. Oppenheim and M. S. Taqqu (eds.) (2003). *Theory and Applications of Long-Range Dependence.* 1st Edition, Birkhäuser.

[9] Durrett, R. (1991). *Probability: Theory and Examples.* Pacific Grove: Wadsworth & Brooks/Cole.

[10] Engle, R.F. (1982). “Autoregressive Conditional Heteroskedasticity with Estimates of Variance of U.K. Inflation”. *Econometrica*, Vol. **50**, 987-1008.

[11] Engle, R.F. and T. Bollerslev (1986). “Modeling the persistence of conditional variances”. *Econometric Reviews*, **5**, 1-50.

[12] Hall, P. and Q. Yao (2003). “Inference in ARCH and GARCH Models with Heavy-tailed Errors”. *Econometrica*, Vol. **71**, 285-317.

[13] Hurvich C.M, E. Moulines and P. Soulier (2005). “ Estimating Long Memory in Volatility”. *Econometrica*, Vol. **73**(4), 1283-1328.

[14] Jayasuriya, S.A. (2009). “A Sub Period Analysis of Long Memory in Stock Return Volatility for Emerging Markets”. *9th Global Conference on Business and Economics Proceedings*, Cambridge University, U.K., 1-31.

[15] Lee, S. and B. Hansen (1994). “Asymptotic Properties of the Maximum Likelihood Estimator and Test of the Stability on the GARCH and IGARCH Models”. *Econometric Theory*, Vol. **10**, 29-52.

[16] Lopes, S.R.C. (2008). “Long-range Dependence in Mean and Volatility: Models, Estimation and Forecasting”, In *In and Out of Equilibrium 2 (Progress in Probability)*, V. Sidoravicius and M.E. Vares (eds.), Vol. **60**, 497-525, Birkhäuser.

[17] Lumsdaine, R. (1996). “Asymptotic Properties of the Maximum Likelihood Estimator in GARCH(1,1) and IGARCH(1,1) Models”. *Econometrica*, Vol. **64**, 575-596.

[18] Mikosch, T. and C. Stărică (1999). “Change of structure in financial time series, long range dependence and the GARCH model“. Preprint.

[19] Ming, L. (2000). “Modeling long memory in stock market volatility”. *Journal of Econometrics*, Vol. **99**(1), 139-171.

[20] Nelson, D.B. (1991). “Conditional Heteroskedasticity in Asset Returns: A New Approach”. *Econometrica*, Vol. **59**, 347-370.

[21] Palma, W. (2007). *Long-Memory Time Series: Theory and Methods.* New Jersey: John Wiley.

[22] Perez, A. and P. Zaffaroni (2008). “Finite-sample Properties of Maximum Likelihood and Whittle Estimators in EGARCH and FIEGARCH Models”. *Quantitative and Qualitative Analysis in Social Sciences*, Vol. **2**(1), 78-97

[23] Prass, T.S. (2008). *Análise e Estimação de Medidas de Risco em Processos FIEGARCH.* Master Dissertation in the Graduate Program in Mathematics, Federal University of Rio Grande do Sul, Porto Alegre.

[24] Prass, T.S. and S.R.C. Lopes (2011a). “Risk Measures Estimation on FIEGARCH Processes”. Working Paper.

[25] Prass, T.S. and S.R.C. Lopes (2011b). “Stress Tests, Maximum Loss and Value-at-Risk on FIEGARCH Process”. Working Paper.

[26] Reisen, V.A. and S.R.C. Lopes (1999). “Some Simulations and Applications of Forecasting Long Memory Time Series Models”. *Journal of Statistical Planning and Inference*, Vol. **80**(2), 269-287.

[27] Ruiz, E. and H. Veiga (2008). “Modelling long-memory volatilities with leverage effect: A-LMSV versus FIEGARCH”. *Computational Statistics and Data Analysis*, Vol. **52**(6), 2846-2862.

[28] Saadi, S., D. Gandhi and S. Dutta (2006). “Testing for nonlinearity and modeling volatility in emerging capital markets: the case of Tunisia”. *International Journal of Theoretical and Applied Finance*, Vol. **9**(7), 1021-1050.

[29] Straumann, D. and T. Mikosch (2006). “Quasi-Maximum-Likelihood Estimation in Conditionally Heteroskedastic Time Series: A Stochastic Recurrence Equations Approach”. *The Annals of Statistics*, Vol. **34**(5), 2449-2495.

[30] Zaffaroni, P. (2005). “Whittle estimation of EGARCH and other exponential volatility models”. *Journal of Econometrics*, Vol. **151** 190-200.

[31] Zivot, E. and J. Wang (2005). *Modeling Financial Time Series with S-PLUS.* New York: Springer-Verlag. 2nd edition.